


**CHEMISTRY MARKING SCHEME**  
**FOREIGN-2013**  
**SET - 56/2/2**

<b>Q no.</b>	<b>Answers</b>	<b>Marks</b>
1	Due to its tendency to flow like liquid	1
2	Ethylamine forms H bond with water but aniline, can't form H bond due to hydrophobic benzene ring	1
3	Phenol < 4-nitrophenol < 2,4,6-trinitrophenol	1
4	$\text{H}_3\text{C}-\text{CO}-\text{CH}=\text{C}(\text{CH}_3)_2$ or structure for m	1
5	Osmotic pressure	1
6	5-chloro-4-methylpent-1-ene	1
7	Differential adsorption	1
8	Ethylene glycol + Terephthalic acid	1
9	Positive deviation Minimum boiling azeotrope	1+1
10	1) Buna-S < Polythene < nylon-6,6 2) Neoprene < Buna-S, nylon-6	1+1
11	<p>Alumina is leached out by using conc. NaOH solution to sodium aluminate and silica as sodium silicate.</p> $\text{Al}_2\text{O}_3 + 2\text{NaOH} + 3\text{H}_2\text{O} \rightarrow 2\text{Na}[\text{Al(OH)}_4]$ <p>Aluminum hydroxide or hydrated alumina is then ppt. by passing CO<sub>2</sub> gas whereas sodium silicate remained in solution.</p> <p>Aluminum hydroxide is ignited to get pure alumina. (or explained in any other correct suitable manner)</p> <p style="text-align: center;">OR</p> <p>(a) <math>\text{Cu}_2\text{S} + \text{FeS}</math> (b) Depressant is used to separate sulphide ore selectively from a mixture of two sulphide ores.</p>	2
11		1
		1


12	<p style="text-align: center;"><i>P</i></p> <p>According to Henry's law, <math>p = k_H x_{\text{CH}_4}</math></p> $\therefore x_{\text{CH}_4} = \frac{p}{k_H} = \frac{760 \text{ mmHg}}{4.27 \times 10^5 \text{ mmHg}} = 1.78 \times 10^{-3}$ <p>Mole fraction of methane in benzene; <math>x_{\text{CH}_4} = 1.78 \times 10^{-3}</math>.</p>	$\frac{1}{2}$ 1 $\frac{1}{2}$
13	<p>a) <math>k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}</math></p> $t = \frac{2.303}{60 \text{ s}^{-1}} \text{ lo}_{\text{o}}$ <p><b>t = 0.0383 sec</b></p>	$\frac{1}{2}$ 1 $\frac{1}{2}$
14	<p>i) <math>\text{CHO}-(\text{CHOH})_4-\text{CH}_2\text{OH} \xrightarrow{\text{HI}} \text{CH}_3-(\text{CH}_2)_4-\text{CH}_3</math></p> <p>ii) <math>\begin{array}{c} \text{CHO} \\   \\ (\text{CHOH})_4 \\   \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{\text{H}_2\text{N-OH}} \begin{array}{c} \text{CH}=\text{N}-\text{OH} \\   \\ (\text{CHOH})_4 \\   \\ \text{CH}_2\text{OH} \end{array}</math></p>	1 1
15	<p>a) Peptization takes place.  b) Because of larger surface area.</p>	1 1
16	<p>(i) Kraft temperature: The temperature above which micellization takes place is called Kraft temperature.</p>	

	(ii) Sorption The phenomenon in which both adsorption and absorption takes place simultaneously.	1+1
17	(i) HF < HCl < HBr < HI (ii) NH <sub>3</sub> < PH <sub>3</sub> < AsH <sub>3</sub> < SbH <sub>3</sub> < BiH <sub>3</sub>	1 1
18	a) Hydrogen bonding b) Nucleotide is sugar +nitrogenous base + phosphate group whereas Nucleoside is sugar + nitrogenous base .	1+1
19	i) Due to discrete tetrahedral structure and angular strain, white phosphorus is more reactive whereas red phosphorus is polymeric and therefore less reactive. ii) Because of higher charge/size ratio of Sn <sup>4+</sup> . iii) Due to its ease of liberating nascent oxygen <b>OR</b>	1x3=3
19	(i) PCl <sub>3</sub> + 3H <sub>2</sub> O → H <sub>3</sub> PO <sub>3</sub> + 3HCl  (ii) XeF <sub>2</sub> + PF <sub>5</sub> → [XeF] <sup>+</sup> [PF <sub>6</sub> ] <sup>-</sup>  (iii) NaN <sub>3</sub> → 2Na + 3N <sub>2</sub>	1x3=3
20	i) Retention of configuration ii) Inversion of configuration iii) Racemisation	1x3=3
21	1) 1 <sup>st</sup> order 2) -k 3) sec <sup>-1</sup>	1x3=3

22



1x3=3




23


- i) Helping, caring and setting an example of true friendship
- ii) Tranquillizers
- iii) Because in excess it acts as poison and can harm the nervous system

1x3=3

24




½



½

1



1



(b) GQ / KMnO<sub>4</sub> / Acidified K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>

25

(a)

- (i)  $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{Cl}]\text{Cl}_2$
- (ii)  $\text{K}_2[\text{N}(\text{CN})_4]$

1+1

	(b) $sp^3$	1
26	$d = \frac{z \times M}{a^3 \times N_A}$ $27 \text{ g cm}^{-3} = \frac{z \times 27 \text{ g mol}^{-1}}{(4.05 \times 10^{-8} \text{ cm})^3 \times 6.022 \times 10^{23} \text{ mol}^{-1}}$ $z = \frac{27 \text{ g cm}^{-3} \times 6.022 \times 10^{23} \text{ mol}^{-1} \times (4.05 \times 10^{-8} \text{ cm})^3}{27 \text{ g mol}^{-1}}$ <div style="border: 1px solid black; padding: 5px; text-align: center;"> <b><math>z \approx 4</math></b> </div>	$\frac{1}{2}$ $1$ $\frac{1}{2}$
	<b>Hence the cubic unit cell is f.c.c.</b>	1
27	(i)  (ii)  (iii) 	1x3=3

28	<p>a)</p> <p>i) Because carbon of carbonyl group in ethanal is more electrophilic than of ketone due to the presence of one electron donating methyl group.  ii) Because of the absence of <math>\alpha</math>-hydrogen atom  iii) Because of extensive association of hydrogen bond / dimerisation in carboxylic acid</p> <p>b)</p> <p>i) Add NaOH + I<sub>2</sub>, acetophenone gives yellow ppt. of CH<sub>3</sub> whereas benzophenone does not form many ppt.  ii) Add NaOH + I<sub>2</sub>, ethanal gives yellow ppt. of CH<sub>3</sub> whereas benzaldehyde does not form many ppt.</p> <p style="text-align: right;"><i>(or any other correct suitable test)</i></p> <p style="text-align: center;">OR</p>	1x3=3
28	<p>i) </p> <p>ii) <math>\text{CH}_3-\overset{\underset{\text{OH}}{\text{ }}}{\text{CH}}-\text{CN}</math></p> <p>iii) <math>\text{HCOO}^-\text{K}^+ + \text{CH}_3\text{OH}</math></p> <p>iv) </p> <p>v) </p>	1 x 5 = 5
29	<p>(a) Kohlrausch's law states that limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the</p>	

	electrolyte	1
	It is used to calculate $\Lambda^0_m$ of even weak electrolyte ./ It is used to calculate degree of dissociation	1
	(b)	
	$R = \rho(l/a)$	1
	<b>Cell constant l/a = R/ρ = Rk</b>	1
	$= 1500 \Omega \times (0.15 \times 10^{-4} \text{ Sc m}^{-1})$	1
	$= 0.225 \text{ cm}^{-1}$	1
29	OR	
	$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$	
	$= 0.34 \text{ V} - (-2.36) \text{ V}$	½
	$= +2.70 \text{ V}$	½
	$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{2} \log \frac{[\text{Mg}^{2+}]}{[\text{Cu}^{2+}]}$	1
	$E_{\text{cell}} = 2.70 \text{ V} - \frac{0.059}{2} \log \left( \frac{0.001 \text{ M}}{0.0001 \text{ M}} \right)$	
	$2.70 \text{ V} - \frac{0.059}{2} \log (10)$	
	$= 2.70 \text{ V} - 0.0295 \text{ V}$	1
	$= 2.6705 \text{ V}$	
	$\Delta G^{\circ} = -nFE_{\text{cell}}^{\circ}$	½
	$= -2 \times 96500 \text{ C mol}^{-1} \times 2.70 \text{ V}$	1
	$= -521.1 \text{ kJ mol}^{-1}$	½

30	<p>i) Because of the absence of unpaired electron in the formation of metallic bond / because of non-involvement of d-orbital electrons in the formation of metallic bond.</p> <p>ii) Because of lanthanoid contraction</p> <p>iii) Because of incomplete filling of d-orbitals.</p> <p>iv) Because of low <math>\Delta_{hyd}</math> H° and high <math>\Delta_a</math> H° of Cu<sup>2+</sup> ion and Cu respectively.</p> <p>v) Because Cr<sup>3+</sup> has stable t<sub>2g</sub><sup>3</sup> half filled configuration</p>	1x5=5
30	<p style="text-align: center;">OR</p> <p>2 MnO<sub>2</sub> + 4KOH + O<sub>2</sub> → 2K<sub>2</sub>MnO<sub>4</sub> + 2H<sub>2</sub>O</p> <p>MnO<sub>4</sub><sup>2-</sup> undergoes disproportionation reaction in acidic medium to give MnO<sub>4</sub><sup>-</sup> ion</p> <p>3 MnO<sub>4</sub><sup>2-</sup> + 4H<sup>+</sup> → 2 MnO<sub>4</sub><sup>-</sup> + MnO<sub>2</sub> + 2H<sub>2</sub>O</p> <p>i) MnO<sub>4</sub><sup>-</sup> + 8H<sup>+</sup> + Fe<sup>2+</sup> → Mn<sup>2+</sup> + Fe<sup>3+</sup> + 4H<sub>2</sub>O</p> <p>ii) 2 MnO<sub>4</sub><sup>-</sup> + 16H<sup>+</sup> + 5C<sub>2</sub>O<sub>4</sub><sup>2-</sup> → 2 Mn<sup>2+</sup> + 10CO<sub>2</sub> + 8H<sub>2</sub>O</p>	1 1 1 1 1
	<p>Sh. S K Murjia</p> <p>Dr (Ms.) Sangeeta Bhatia</p> <p>Prof. R D Shukla</p> <p>Dr. K N Uppadhyaya</p> <p>Mr. Rakesh Dhawan</p> <p>Ms. Neeru Sofat</p> <p>Mr. Virendra Singh</p> <p>M. K M Abdul Raheem</p> <p>M. D A Mishra</p> <p>M. Deshbir Singh</p> <p>M. Akhileshwar Mishra</p>	