


CHEMISTRY MARKING SCHEME


Bhubaneswar – 2015

Set 2 - Code No. 56/2/B

Ques.	Value points	Marks
1.	HOCl , HOClO, HOClO ₂ , HOClO ₃ (Any two of these)	½ +½
2.	$\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{CH}_2 - \text{Br} \\ \\ \text{CH}_3 \end{array}$	1
3.	Negative charge	1
4.	XY ₃	1
5.	1-Phenylpropan-2-ol	1
6.	<p>Formula: $w = z \times i \times t$</p> $\text{time taken in sec} = \frac{w \times \text{Valance} \times 96500}{\text{Mol Mass} \times \text{Current in Amp}}$ <p>Substituting the values in the formula we get:</p> $\text{time taken in sec} = \frac{1.17 \text{ g} \times 2 \times 96500 \text{ C mol}^{-1}}{58.5 \text{ g mol}^{-1} \times 5 \text{ amp}}$ $\text{time taken in sec} = \frac{225810}{292.5}$ <p style="text-align: center;">t=772 s</p> <p>(Or by any other correct method)</p>	<p>½</p> <p>1</p> <p>½</p>
7.	(i) Potassium hexacyanidoferrate (III)	1
	(ii) [Co(NH ₃) ₅ NO ₂] ²⁺	1
8.	(i) Due to comparable energies of 5f, 6d and 7s orbitals .	1
	(ii) Because 5f electrons have poorer shielding effect than 4f electrons.	1
9.	(i) Positive deviation, lowering of temperature or absorption of heat. (ii) By applying an external pressure greater than the osmotic pressure on the solution or $P > \pi$ Reverse osmosis is used in desalination of hard water / sea water.	<p>½ ,½</p> <p>½ , ½</p>
10.	(i) H ₂ / Pd-BaSO ₄ (ii) NaOH/CaO, Δ	1 1
	OR	
10.	i) C ₆ H ₅ CO C ₆ H ₅ < CH ₃ COCH ₃ < CH ₃ CHO	1
	ii) Cl – CH ₂ – COOH < Cl ₂ CH – COOH < CCl ₃ – COOH	1

11.	(i) Distillation (ii) Collector/ enhancing the non-wettability of mineral particles. (iii) As ΔS is positive / ΔG is more negative	1 1 1
12.	(i) Stoichiometric Defect (ii) Frenkel Defect (iii) Due to small size of Ag^+ ion	1 1 1
13.	(i) $CH_3 - CH(OH) - CN$ (ii) $C_6H_5 - COOH$ (iii) $CH_3 - CH_2NH_2$	1 1 1
14.	(ii) $t_{2g}^3 e_g^1$ (iii) Hybridization dsp^2 , Shape \rightarrow Square planar or diagram <p>(Marks of (i) part is merged into (ii) and (iii) part)</p>	1 ½ 1 ½
15.	(i) Due to the stability of benzyl carbocation/resonance/Diagram (ii) Because 2-Bromobutane has a chiral centre. (iii) Due to - I effect of halogen.	1 1 1
16.	(i) $C_6H_5NH_2 \xrightarrow[0^\circ - 5^\circ C]{NaNO_2 + HCl} C_6H_5N_2Cl \xrightarrow[Or Hydrolysis]{H_2O + H^+} C_6H_5OH$ (ii) $CH_3 - CH = CH_2 \xrightarrow[Organic peroxide]{HBr} CH_3 - CH_2 - CH_2Br \xrightarrow{KOH_{Aq}} CH_3CH_2CH_2OH$ (iii) <p style="text-align: right;">(Or any correct method)</p>	1 1 1

	<p>1,1,2,2-Tetrafluoroethene</p> <p>(iii) Nylon-6 Monomer: Caprolactum</p>  <p>Caprolactum</p> <p>(Note : half mark for structure/s and half mark for name/s)</p>	1
20.	<p>(i) Because of higher oxidation state of Mn in Mn_2O_7.</p> <p>(ii) Due to almost similar atomic size / comparable size.</p> <p>(iii) $2MnO_2 + 4KOH + O_2 \longrightarrow 2K_2MnO_4 + 2H_2O$</p>	1 1 1
21.	<p>(i) Solution is homogeneous colloid is heterogeneous In solution the size of particles (solute) is less than 1 nm whereas in colloids the range of size of particles is 1 – 1000 nm (10^{-9} to 10^{-6} m)(Any one point)</p> <p>(ii) In homogeneous catalysis the reactant and catalyst are in the same phase whereas in heterogeneous catalysis they are in different phase.</p> <p>(iii) In O/W emulsion oil is the dispersed phase while in W/O water is dispersed in oil The O/W type emulsion can be diluted with water whereas the W/O emulsion can't be diluted with water. (Any one point)</p>	1 1 1
22.	<p>Formula $\frac{p_1^0 - p_1}{p_1^0} = \frac{w_2 \times M_1}{M_2 \times w_1}$</p> $\frac{23.75 \text{ mm} - 23.375 \text{ mm}}{23.75 \text{ mm}} = \frac{5.0 \text{ g} \times 18 \text{ g/mol}}{M_2 \times 95.0 \text{ g}}$ $M_2 = \frac{5.0 \text{ g} \times 18.0 \text{ g/mol} \times 23.75 \text{ mm}}{95 \text{ g} \times 0.375 \text{ mm}}$ <p>$M_2 = 60.0 \text{ g/mol}$</p>	1 1 1
23.	<p>(i) Concern for students health, Application of knowledge of chemistry to daily life, empathy , caring or any other (Any two)</p> <p>(ii) Through posters, nukkad natak in community, social media, play in assembly or any other (Any two)</p> <p>(iii) Tranquilizers are drugs used for treatment of stress or mild and severe mental disorders . Eg: equanil (or any other suitable example)</p> <p>(iv) Aspartame is unstable at cooking temperature.</p>	$\frac{1}{2}$, $\frac{1}{2}$ 1 $\frac{1}{2}$, $\frac{1}{2}$ 1
24.	<p>a)</p> <p>(i) The +3 Oxidation state of Bi is more stable than Sb(III) .</p> <p>(ii) Because the electronegativity of Cl is greater than that of I .</p> <p>(iii) Due to decrease in electronegativity and increase in the atomic size.</p>	1 1 1

25.	 <p>b) $(\text{CH}_3)_3\text{N} < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$</p> <p>c) Dye Test: On treating with benzene diazonium Chloride at low temperature $\text{C}_6\text{H}_5\text{-NH}_2$ will form coloured dye while $\text{CH}_3\text{-NH}_2$ will not form. (or any other correct distinguishing test)</p>	1 1 1 1 1
26.	<p>(a)</p> <p>Formula: $k = \frac{2.303}{t} \log \frac{[\text{CH}_3\text{COOCH}_3]_1}{[\text{CH}_3\text{COOCH}_3]_2}$</p> $k_1 = \frac{2.303}{20\text{s}} \log \frac{0.4\text{M}}{0.2\text{M}}$ $k_1 = 0.03 \text{ s}^{-1}$ $k_2 = \frac{2.303}{40\text{s}} \log \frac{0.4\text{M}}{0.1\text{M}}$ $k_2 = 0.03 \text{ s}^{-1}$ <p>Since constant values of rate constants are obtained by applying 1st Order integrated rate law, the reaction is pseudo first order reaction.</p> <p>(b) $\text{Av rate} = \frac{\text{total change in concentration}}{\text{total change in time}}$</p> <p>OR</p> $\text{Av rate} = - \frac{[\text{CH}_3\text{COOCH}_3]_{\text{final}} - [\text{CH}_3\text{COOCH}_3]_{\text{initial}}}{\text{Time}(f) - \text{Time}(i)}$ $\text{Av rate} = - \frac{0.10\text{M} - 0.20\text{M}}{40\text{Sec} - 20\text{Sec}}$ $\text{Av rate} = 0.0005 \text{ M sec}^{-1} \text{ or } 5.0 \times 10^{-3} \text{ mol L}^{-1} \text{ sec}^{-1}$ <p>OR</p> <p>a) i) Collision frequency: No of collisions taking place per second per unit volume. ii) Rate Constant: It is the rate of reaction when the concentration of reactants</p>	½ 1 1 ½ ½ 1 ½ 1
26.		1

	is unity i.e. 1 M. It is temperature dependent	1
b)	$\log \frac{k_2}{k_1} = \frac{Ea}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$	1
	$\log \frac{k_2}{k_1} = \frac{Ea}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$	
	$\log 6 = \frac{Ea}{19.147} \left[\frac{50}{105000} \right]$	1
	$0.7782 = \frac{Ea}{19.147} \left[\frac{50}{105000} \right]$	
	$0.7782 = \frac{Ea}{19.147} [0.00047619]$	
	$\frac{0.7782 \times 19.147}{0.00047619} = Ea = 31290.44 \text{ J}$	1
	$Ea = 31.29 \text{ kJ/mol}$	