










MARKING SCHEME CHEMISTRY-2015**(CODE NO. : 56/3/C)**




Q	Value points	Mark						
1	2-Methylpropane-1,3-diol	1						
2	White phosphorous, because of angular strain in P ₄ molecule/ discrete tetrahedral unit.	1						
3	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{Br} \\ \\ \text{CH}_3 \end{array}$ <p>Because carbocation intermediate derived from (CH₃)₃Br is more stable than carbocation from CH₃CH₂Br.</p>	½						
4	(i) Electrophoresis (ii) by mixing two oppositely charged sols (iii) by boiling (iv) by persistent dialysis (v) by addition of electrolyte (any one)	1						
5	X ₄ Y ₃	1						
6	(i) pentaamminenitrito-N-cobalt(III) nitrate (ii) K ₂ [Ni(CN) ₄]	1 1						
7	(i) CH ₃ MgBr, H ₃ O ⁺ (ii) Cl ₂ , P	1 1						
8	It states that solubility of gas in liquid is directly proportional to partial pressure of the gas in equilibrium with the solution. With increase in temperature K _H value increases but solubility of gas in liquid decreases. / K _H ∝ 1/solubility OR 8. It states for solution containing volatile components the partial vapor pressure of each component of the solution is directly proportional to its mole fraction present in the solution.	1 1 1						
	<table border="1"> <thead> <tr> <th>Ideal Solution</th> <th>Non Ideal</th> </tr> </thead> <tbody> <tr> <td>1. It obeys Raoult's Law over entire range of concentration of solution.</td> <td>It does not obey Raoult's Law.</td> </tr> <tr> <td>2. Solute – Solvent interaction is nearly same as in pure solvent.</td> <td>Solute – Solvent interaction is not same as solute-solute or solvent – solvent interactions.</td> </tr> </tbody> </table>	Ideal Solution	Non Ideal	1. It obeys Raoult's Law over entire range of concentration of solution.	It does not obey Raoult's Law.	2. Solute – Solvent interaction is nearly same as in pure solvent.	Solute – Solvent interaction is not same as solute-solute or solvent – solvent interactions.	½ + ½
Ideal Solution	Non Ideal							
1. It obeys Raoult's Law over entire range of concentration of solution.	It does not obey Raoult's Law.							
2. Solute – Solvent interaction is nearly same as in pure solvent.	Solute – Solvent interaction is not same as solute-solute or solvent – solvent interactions.							
	(or any other correct difference)							
9	Greater number of unpaired electron, greater is the interatomic interactions leading to strong metallic bonding.	1						

	Zn , no unpaired electrons hence weak metallic bonding.	1/2 , 1/2
10	(a) $H^+ (aq) + e^- \rightarrow 1/2H_2 (g)$ $E^\circ = 0.00 V$ is feasible at cathode because its reduction potential is higher than the other reaction. b. Because the overall reaction doesn't involve any ion in the solution whose concentration changes during its lifetime.	1/2 , 1/2 1
11	(i) $CH_3-CH=CH_2 \xrightarrow[\text{peroxide}]{HBr} CH_3-CH_2-CH_2-Br \xrightarrow{AgF} CH_3CH_2CH_2F$ (ii)  (iii) $C_2H_5OH \xrightarrow{PCl_3/PCl_5} C_2H_5Cl \xrightarrow{KCN} C_2H_5CN$ OR (i) $CH_3CH_2CH=CH_2$ (ii)  (iii) CH_3NC	1 1 1 1,1,1
12	(i) Because $-NO_2$ is an electron withdrawing group (ii) Due to H-Bonding (iii) Reaction occurs by S_N1 mechanism 3° -carbocation $(CH_3)_3C^+$ is more stable than CH_3^+	1 1 1
13	$\Delta T_f = i \times K_f \times m$ For $CaCl_2$ $i = 3$	1/2 1/2

	$\Delta T_f = (i \times K_f \times W_B \times 1000) / (M_B \times W_A)$ $2 = 3 \times 1.86 \times W_B \times 1000 / 111 \times 500$ $W_B = 19.89 \text{ g}$	1 1
14	$d = Z \times M / a^3 \times N_o$ $10 \text{ g/cm}^3 = Z \times 81 \text{ g/mol} / (3 \times 10^{-8} \text{ cm})^3 \times (6.023 \times 10^{23} / \text{mol})$ $Z = 2.007$ <p>Nature of cubic unit cell = bcc</p>	1/2 1/2 1 1
15	$E^\circ_{\text{cell}} = E_R^\circ - E_L^\circ$ $= 0.00 - (-0.14)$ $E^\circ_{\text{cell}} = +0.14 \text{ V}$ $E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.059 \text{ V}}{n} \log \frac{[\text{Sn}^{2+}]}{[\text{H}^+]^2}$ $E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.059 \text{ V}}{2} \log \frac{[0.001]}{[0.01]^2}$ $= +0.14 - 0.0295 \text{ V} \log 10$ $E_{\text{cell}} = 0.1105 \text{ V}$	1 1 1
16	 <p>(i) cis</p>  <p>(ii) trans</p> <p>(iii) $[\text{NiCl}_4]^{2-}$ -Chloride ion being weak field ligand does not pair d electrons while in $[\text{Ni}(\text{CO})_4]$, CO being strong field ligand pairs up the d electrons.</p>	1 1 1
17	<p>(i) Because physisorption is exothermic process, so it decreases with increase in temperature.</p> <p>(ii) Because alum coagulates the impurities present in water.</p> <p>(iii) Due to continuous unbalanced bombardment / zig-zag motion of particles by the molecules of dispersion medium / it allows the particles to settle down.</p>	1 1 1
18	<p>(i) van Arkel method</p> <p>(ii) CO acts as reducing agent</p> <p>(iii) Because ΔS becomes more positive, and ΔG becomes negative.</p>	1 1 1

19	<div style="text-align: center;">  <p>(i) Caprolactum Caprolactum</p> </div> <p>(ii) Phenol + Formaldehyde</p> <div style="text-align: center;">  <p>+ HCHO</p> </div> <p>(iii) 1,3-Butadiene + Acrylonitrile $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + \text{CH}_2=\text{CH}-\text{CN}$ (Note: half mark for structure/s and half mark for name/s)</p>	<p>1</p> <p>1</p> <p>1</p>
20	<p>(i) Starch</p> <p>(ii) Native Protein found in a biological system with a unique 3-D structure and biological activity is called a native protein. Denatured protein is the protein with no biological activity.</p> <p>(iii) Vitamin-K</p>	<p>1</p> <p>1</p> <p>1</p>
21	<p>(i) $\text{CH}_3 - \text{CH}(\text{OH}) - \text{CN}$</p> <p>(ii) $\text{C}_6\text{H}_5\text{COOH}$</p> <p>(iii) CH_3CONH_2</p>	<p>1</p> <p>1</p> <p>1</p>
22	<p>(a)(i) Because actinoids are radioactive and show wide range of oxidation states.</p> <p>(ii) Transition metals form complex compounds due to - small size, high ionic charge, availability of d orbitals</p> <p>b. $2\text{MnO}_4^- + 6\text{H}^+ + 5 \text{SO}_3^{2-} \rightarrow 5\text{SO}_4^{2-} + 3\text{H}_2\text{O} + 2 \text{Mn}^{2+}$</p>	<p>1</p> <p>1</p> <p>1</p>
23	<p>(i) Concern, Compassion, caring, empathy (any two)</p> <p>(ii) By organizing rallies, street play, posters, public speech (any other relevant answer)</p> <p>(iii) Anti depressant drugs are those which inhibit depression E.g. Iproniazide, Phenelzine (or any other)</p> <p>(iv) Saccharine / Sucralose / Alitame / Aspartame (any one)</p>	<p>$\frac{1}{2}, \frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}, \frac{1}{2}$</p> <p>1</p>
24	<p>(i) X-X' bond in inter halogens is weaker than X-X in halogens</p> <p>(ii) High bond dissociation energy/ due to presence of triple bond.</p> <p>(iii) Because bond dissociation enthalpy decreases from NH_3 to BiH_3.</p>	<p>1</p> <p>1</p> <p>1</p>

24	<p>b. (i)</p>  <p>(ii)</p>  <p>OR</p> <p>a) PH_3 $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow 3\text{NaH}_2\text{PO}_2 + \text{PH}_3$</p> <p>b) Helium</p> <p>c) Because bond dissociation energy of F-F bond is lower than that of Cl-Cl.</p> <p>d) $4\text{H}_3\text{PO}_3 \xrightarrow{\text{HEAT}} 3\text{H}_3\text{PO}_4 + \text{PH}_3$</p> <p>e) $\text{PbS} + 4\text{O}_3 \rightarrow \text{PbSO}_4 + 4\text{O}_2$</p>	1,1 1/2, 1/2 1 1 1 1
25	<p>(i) $\text{CH}_3\text{CONH}_2 + \text{Br}_2 + 4\text{KOH} \rightarrow \text{CH}_3\text{NH}_2 + \text{K}_2\text{CO}_3 + 2\text{KBr} + 2\text{H}_2\text{O}$</p> <p>(ii) $\text{C}_6\text{H}_5\text{NH}_2 + \text{NaNO}_2 + 2\text{HCl} \xrightarrow{273-278\text{K}} \text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^- + \text{NaCl} + 2\text{H}_2\text{O}$</p> <p>(iii)</p>  <p>b.(i) Add CHCl_3 and alc KOH, aniline gives foul smell of isocyanide whereas N-methylaniline does not.</p> <p>(ii) When $(\text{CH}_3)_2\text{NH}$ reacts with Benzene Sulphonyl Chloride (Hinsberg Reagent) gives ppt which is insoluble in alkali whereas $(\text{CH}_3)_3\text{N}$ does not react with Hinsberg's Reagent.</p> <p>(Or any other correct distinguishing test)</p> <p>OR</p>	1 1 1 1 1

25	<p>a.</p> <p>(i) </p> <p>(ii) </p> <p>(iii) </p> <p>b. (i) $(\text{CH}_3)_3\text{N} < \text{C}_2\text{H}_5\text{NH}_2 < \text{C}_2\text{H}_5\text{OH}$ (ii) p-nitroaniline < aniline < p-methylaniline</p>	1,1,1 1 1
26	<p>a. Rate of reaction is defined as change in concentration of reactants or products per unit time. Factors: concentration of reactant , temperature , pressure , surface area (any two)</p> <p>b. $\log(k_2/k_1) = E_a/2.303R [1/T_1 - 1/T_2]$</p> <p>$\log(8 \times 10^{-2}/4 \times 10^{-2}) = E_a/2.303 \times 8.314 [1/300 - 1/310]$</p> <p>$\log 2 = E_a/2.303 \times 8.314 [1/300 - 1/310]$ $E_a = 53598.59 \text{ J/mol or } 53.6 \text{ kJ/mol}$</p> <p style="text-align: center;">OR</p>	1 1/2, 1/2 1 1 1
26	<p>(a)(i) Rate becomes 4 times (ii) 2nd order</p> <p>b) $t_{1/2} = \frac{0.693}{k}$</p> <p>$23.1 \text{ min} = \frac{0.693}{k}$</p> <p>$k = 0.03 \text{ min}^{-1}$</p> <p>$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$</p> <p>$t = \frac{2.303}{0.03} \log \frac{100}{25}$</p>	1 1 1 1/2 1/2

	$t = \frac{2.303}{0.03} \times 0.6021 \text{ min}$ $t = 46.22 \text{ min}$	1
--	--	---