











CHEMISTRY MARKING SCHEME 2015**PATNA****SET -56/1/P**

Qu es.	Value points	Marks
1	2F or 2x 96500C	1
2	Dispersed phase -liquid Dispersion medium - solid	½ +½
3	Because of no unpaired electron in Zn^{2+} Copper salts are coloured due to the presence of unpaired electrons in Cu^{2+}	½ +½
4	2-Methyl prop-2-en-1-ol	1
5	$(\text{CH}_3)_3\text{C-Br}$	1
6.	Because on addition of a non- volatile solute, vapour pressure of solution lowers down and therefore in order to boil solution, temperature has to be increased, thus boiling point gets higher Because it depends on molality/ number of solute particles / $\Delta T_b \propto m$	1 1
7.	Decrease in concentration of reactant or increase in concentration of product per unit time Factrors: 1)concentration of reactant 2)catalyst 3) temperature 4)Nature of reactant 5)pressure 6)surface area (any two)	1 ½ +½
8.	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(i)</p> </div> <div style="text-align: center;">  <p>(ii)</p> </div> </div>	1,1
9	Dichloridobis-(ethane-1,2-diamine)platinum(IV) Geometrical or optical isomerism OR	1 1
9.	(i) $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ (ii) $\text{K}_2[\text{NiCl}_4]$	1 1
10	(i) $\text{C}_6\text{H}_5\text{NH}_2 < \text{C}_6\text{H}_5\text{NHCH}_3 < \text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$	1

	<p>(ii)</p> 	1
11	$\Delta T_f = K_f m$ $T_f^0 - T_f = \frac{K_f W_B \times 1000}{M_B \times W_A}$ $273K - T_f = 1.86K \text{ kg mol}^{-1} \times \frac{31g}{62g\text{mol}^{-1}} \times \frac{1000}{500kg}$ $T_f = (273 - 1.86) K$ $T_f = 271.14K \quad \text{Or} \quad -1.86^\circ C$	1 1 1
12	<p>(i) Unit cells having constituent particles at the corner positions.</p> <p>(ii) The defect occurs due to missing of equal no of cations and anions in a lattice.</p> <p>(iii) The permanent magnetism which arises when magnetic moments of substance are aligned in same direction.</p>	1 1 1
13	$\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$ $\log \frac{4 \times 10^{-2}}{2 \times 10^{-2}} = \frac{E_a}{2.303 \times 8.314 J/K/mol} \left[\frac{1}{300} - \frac{1}{310} \right]$ $\log 2 = \frac{E_a}{19.147 J/mol} \left[\frac{10}{300 \times 310} \right]$ $E_a = \frac{0.3010 \times 19.147 \times 300 \times 310}{10}$ $E_a = 53598 J/mol \quad \text{or} \quad 53.598 kJ/mol$	1 1 1
14	<p>(i) The zig-zag motion of the colloidal particles due to unbalanced bombardment by the particles of dispersion medium.</p> <p>(ii) The conversion of precipitate into colloidal sol by adding small amount of an electrolyte.</p> <p>(iii) On dissolution a large number of atoms or smaller molecules of a substance aggregate together to form species having size in the colloidal range.</p>	1 1 1
15	<p>(i) Greater solubility of impurities in molten state.</p> <p>(ii) Silica reacts with impurity FeO to form slag (FeSiO_3) / acts as a flux to remove impurities.</p> <p>(iii) Cast iron is harder than pig iron / has lesser content of carbon.</p>	1 1 1
16	<p>(i) Because of the presence of triple bond between two N atoms / high bond dissociation enthalpy</p> <p>(ii) Because of the lowest bond dissociation enthalpy / least thermal stability.</p> <p>(iii) Because of low solubility in blood.</p>	1 1 1
17	<p>(i) $[\text{CoF}_6]^{3-}$ sp^3d^2, octahedral</p>	$\frac{1}{2}$ $\frac{1}{2}$

	(ii) $[\text{Ni}(\text{CN})_4]^{2-}$ dsp^2 , square planar	$\frac{1}{2}$ $\frac{1}{2}$
	(b) CO, because of synergic /back bonding with metal	$\frac{1}{2}$ $\frac{1}{2}$
18	<p>i) $\text{CH}_3 - \text{CH}_2 - \overset{\text{Br}}{\underset{\text{CH}_3}{\text{C}}} - \text{CH}_3$</p> <p>ii) $\text{CH}_3 - \text{CH}_2 - \text{CH} = \text{CH} - \text{CH}_3$</p> <p>iii) </p>	<p>1</p> <p>1</p> <p>1</p>
19	<p>(i) Because phenoxide ion is more stable than $\text{CH}_3\text{CH}_2\text{O}^-$ ion / due to resonance in phenol, oxygen acquires positive charge and releases H^+ ion easily whereas there is no resonance in $\text{CH}_3\text{CH}_2\text{OH}$</p> <p>(ii) Because of hydrogen bonding in ethanol</p> <p>(iii) Because it follows $\text{S}_{\text{N}}1$ path way which results in the formation of stable $(\text{CH}_3)_3\text{C}^+$.</p>	<p>1</p> <p>1</p> <p>1</p>
20	<p>(i) $\text{C}_6\text{H}_5\text{CONH}_2 \xrightarrow{\text{Br}_2 + \text{KOH}} \text{C}_6\text{H}_5\text{NH}_2$</p> <p>(ii) $\text{C}_6\text{H}_5\text{NH}_2 \xrightarrow[0-5^\circ\text{C}]{\text{NaNO}_2 + \text{HCl}} \text{C}_6\text{H}_5\text{N}^+\text{H}_2\text{Cl}^- \xrightarrow{\text{H}_2\text{O}} \text{C}_6\text{H}_5\text{OH}$</p> <p>(iii) $\text{CH}_3\text{CN} \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CH}_2\text{NH}_2$</p> <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p> <p>1</p>

20.	<p>(i)</p> <div style="text-align: center;">  </div> <p>(ii)</p> $\text{R-NH}_2 + \text{CHCl}_3 + 3\text{KOH} \xrightarrow{\text{Heat}} \text{R-NC} + 3\text{KCl} + 3\text{H}_2\text{O} \quad (\text{R} = -\text{C}_6\text{H}_5)$ <p>(iii)</p> $\text{C}_6\text{H}_5\text{NH}_2 + \text{HCl} \longrightarrow \text{C}_6\text{H}_5\text{NH}_3^+ \text{Cl}^-$	<p>1</p> <p>1</p> <p>1</p>									
21	<table border="0" style="width: 100%;"> <tr> <td style="vertical-align: top;">i)Buna –S</td><td style="vertical-align: top;">Butadiene CH₂=CH–CH=CH₂</td><td style="vertical-align: top;">Styrene C₆H₅CH=CH₂.</td></tr> <tr> <td style="vertical-align: top;">ii)Glyptal</td><td style="vertical-align: top;">Ethylene Glycol HO–CH₂CH₂–OH</td><td style="vertical-align: top;">Pthalic acid </td></tr> <tr> <td style="vertical-align: top;">iii)Polyvinyl chloride</td><td style="vertical-align: top;">Vinyl Chloride CH₂=CH–Cl</td><td></td></tr> </table> <p style="text-align: right;">(Note: half mark for name/s and half mark for structure/s)</p>	i)Buna –S	Butadiene CH₂=CH–CH=CH₂	Styrene C ₆ H ₅ CH=CH ₂ .	ii)Glyptal	Ethylene Glycol HO–CH₂CH₂–OH	Pthalic acid 	iii)Polyvinyl chloride	Vinyl Chloride CH₂=CH–Cl		<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>½ ½</p>
i)Buna –S	Butadiene CH₂=CH–CH=CH₂	Styrene C ₆ H ₅ CH=CH ₂ .									
ii)Glyptal	Ethylene Glycol HO–CH₂CH₂–OH	Pthalic acid 									
iii)Polyvinyl chloride	Vinyl Chloride CH₂=CH–Cl										
22	<p>i)</p> <div style="text-align: center;">  </div> <p>(ii)Because of zwitter ion nature of amino acid /</p> <p>(iii)Because vitamin C is soluble in water.</p> <div style="text-align: center;">  </div>	<p>1</p> <p>1</p> <p>1</p>									

23	<p>i) Caring ,concerned, helping,empathy (any two)</p> <p>ii) By organizing competitions like slogan writing, poster making and talk in the morning assembly (any other correct answer)</p> <p>iii) Used to treat depression,. Iproniazid/phenelzine (any other correct example)</p> <p>iv) Saccharin/ sucralose/aspartame/alitame (any other correct example)</p>	<p>$\frac{1}{2}$ $\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$ $\frac{1}{2}$</p> <p>1</p>
24	<p>a) $\text{CH}_3\text{CO Cl}$ (A) CH_3CHO (B) $\begin{array}{c} \text{OH} \\ \\ \text{CH}_3\text{CH}-\text{CH}_2-\text{CHO} \end{array}$ (C) $\text{CH}_3\text{CH}=\text{CH}-\text{CHO}$ (D)</p> <p>b) i)On adding Tollen's reagent $\text{C}_6\text{H}_5\text{CHO}$ forms silver mirror whereas $\text{C}_6\text{H}_5\text{COCH}_3$ does not.</p> <p>ii)On adding NaHCO_3 solution benzoic acid gives brisk effervescence but methyl benzoate does not.</p> <p>(or any other distinguishing test)</p> <p>c) $\begin{array}{c} \text{CH}_3\text{CH}_2-\text{CH}-\text{CHO} \\ \\ \text{CH}_3 \end{array}$</p> <p style="text-align: center;">OR</p> <p>a)i) $\text{CH}_3\text{CH}_2\text{CH}_3$</p> <p>ii) $\begin{array}{c} \text{CH}_3-\text{C}=\text{N}-\text{NHCONH}_2 \\ \\ \text{CH}_3 \end{array}$</p> <p>iii) $\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3-\text{C}-\text{OH} \\ \\ \text{CH}_3 \end{array}$</p> <p>b) $\text{CH}_3\text{CHO} < \text{CH}_3\text{CH}_2\text{OH} < \text{CH}_3\text{COOH}$</p> <p>c)On adding Tollen's reagent $\text{CH}_3\text{CH}_2\text{CHO}$ forms silver mirror whereas $\text{CH}_3\text{CH}_2\text{COCH}_3$ does not (or any other distinguishing test).</p>	<p>$\frac{1}{2}$,$\frac{1}{2}$</p> <p>$\frac{1}{2}$,$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
24		

25	<p> $\text{Mg} \mid \text{Mg}^{2+} (0.001) \parallel \text{Cu}^{2+} (0.0001\text{M}) \mid \text{Cu}$ $E_{\text{Cell}}^0 = E_{\text{R}}^0 - E_{\text{L}}^0$ $= [0.34 - (-2.37)] \text{ V}$ $= 2.71 \text{ V}$ $E_{\text{cell}} = E_{\text{Cell}}^0 - \frac{0.059}{n} \text{ V} \log \frac{[\text{Mg}^{2+}]}{[\text{Cu}^{2+}]}$ $= 2.71 \text{ V} - \frac{0.059}{2} \text{ V} \log 10^{-3}/10^{-4}$ $= 2.71 - 0.0295 \text{ V} \log 10$ $= 2.71 - 0.0295$ $= 2.6805 \text{ V}$ $\Delta G = -nFE_{\text{cell}}$ $= -2 \times 96500 \text{ C mol}^{-1} \times 2.68 \text{ V}$ $= -517240 \text{ J mol}^{-1}$ $= \mathbf{-517.240 \text{ kJ/mol}}$ </p> <p style="text-align: center;">OR</p>	1 1 1 1/2 1/2 1
25.	<p> a) M=0.20M K = 2.48X10⁻²S/cm $\Lambda_m = \frac{K}{M} \times 1000 \text{ Scm}^2/\text{mol}$ $\Lambda_m = \frac{2.48 \times 10^{-2}}{0.20} \times 1000 \text{ Scm}^2/\text{mol}$ $= 124 \text{ Scm}^2/\text{mol}$ $\alpha = \frac{\Lambda_m}{\Lambda_m^0}$ $\Lambda_m^0 = \lambda^0 K^+ + \lambda Cl^-$ $= 73.5 + 76.5$ $= 150$ $\alpha = \frac{124}{150} = 0.82 \quad \text{Or} \quad 82\%$ </p> <p>b) Primary battery or cell, potential remains constant throughout its life.</p>	1/2 1 1/2 1 1,1

26	<p>a)</p> <p>i) Due to lanthanoid contraction.</p> <p>ii) Due to incomplete filling of d- orbitals / comparable energies of (n-1)d & ns electrons.</p> <p>iii) Because it undergoes disproportionation reaction in aqueous solution/ oxidation of a metal in a solvent depends on the nature of the solvent. Cu^+ is unstable in water that's why it undergoes oxidation.</p> <p>b)</p> <p>i) $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \rightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$</p> <p>ii) $2\text{Na}_2\text{CrO}_4 + 2\text{H}^+ \rightarrow \text{Na}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{O} + 2\text{Na}^+$</p> <p style="text-align: center;">OR</p>	1 1 1 1 1
26.	<p>a)</p> <p>(i) Because of high $\Delta_a H^\circ$ & low $\Delta_{\text{hyd}} H^\circ$.</p> <p>(ii) Because of more stability of Mn^{2+} ($3d^5$)</p> <p>(iii) Cr^{2+}, because in +3 oxidation state Cr is more stable (t_{2g}^3 orbital)</p> <p>b) Due to comparable energies of 5f, 6d, 7s orbitals.</p> <p>Both show contraction in size/ both show main oxidation state +3/both are electro positive and very reactive/ both exhibit magnetic and spectral properties. (any one)</p>	1 1 $\frac{1}{2}, \frac{1}{2}$ 1 1