







Marking scheme – 2017**CHEMISTRY (043)/ CLASS XII****FOREIGN 2017 - Set - 56/2/1**

Q.NO.	VALUE POINTS	MARKS
1	P ₃ Q ₄	1
2	H ₂ Te < H ₂ Se < H ₂ S < H ₂ O	1
3	To make the surface available again for more reaction to occur / To remove the product formed from the surface of the catalyst.	1
4	2 – Phenylethanol	1
5	Neopentane / C(CH ₃) ₄	1
6	<p>a.</p> $\text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow[\text{H}^+]{\text{H}_2\text{O}} \text{CH}_3\text{CH}(\text{OH})\text{CH}_3 \xrightarrow{\text{CrO}_3} \text{CH}_3\text{COCH}_3$ <p>b.</p> $\text{CH}_3\text{CH}_2\text{COOH} \xrightarrow{\text{Br}_2/\text{Red P}} \text{CH}_3\text{CH}(\text{Br})\text{COOH} \xrightarrow[\text{ii) H}^+]{\text{i) aq KOH or NaOH}} \text{CH}_3\text{CH}(\text{OH})\text{COOH}$ <p style="text-align: center;">(or any other suitable method)</p>	1
	OR	
6	<p>a. Etard reaction:</p> $\text{Toluene} + \text{CrO}_2\text{Cl}_2 \xrightarrow{\text{CS}_2} \text{Chromium complex} \xrightarrow{\text{H}_3\text{O}^+} \text{Benzaldehyde}$ <p style="text-align: center;">or</p> $\text{Toluene} \xrightarrow[\text{(ii) H}^+\text{O}^+]{\text{(i) CrO}_2\text{Cl}_2, \text{CS}_2} \text{Benzaldehyde}$ <p>b. Wolff-Kishner reduction:</p> $\text{C=O} \xrightarrow{-\text{H}_2\text{O}} \text{C}=\text{NNH}_2 \xrightarrow[\text{heat}]{\text{KOH/ethylene glycol}} \text{CH}_2 + \text{N}_2$ <p style="text-align: center;">or</p> $\text{C=O} \xrightarrow[\text{(ii) KOH/ethylene glycol, heat}]{\text{(i) NH}_2\text{NH}_2} \text{CH}_2 + \text{N}_2$	1

7	Properties that depend on the number of solute particles irrespective of their nature relative to the total number of particles present in the solution. Osmotic Pressure	1
8	a. cis/ trans-diamminedichloroplatinum(II) b. $[Co(NH_3)_4(H_2O)Cl](NO_3)_2$	1 1
9	a. Zinc to silver b. Concentration of Zn^{2+} ions will increase and Ag^+ ions will decrease.	1 1
10	a. Cr^{3+} b. Mn^{3+} c. Ti^{4+} d. Mn^{3+}	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
11	$A = \pi r^2$ $= 3.14 \times 0.5 \times 0.5 \text{ cm}^2$ $= 0.785 \text{ cm}^2$ $l = 45.5 \text{ cm}$ $\rho = R \times A / l$ $\rho = 4.55 \times 10^3 \Omega \times 0.785 \text{ cm}^2 / 45.5 \text{ cm}$ $\rho = 78.5 \Omega \text{ cm}$ conductivity, $\kappa = 1/\rho$ $= 1/78.5 \text{ S cm}^{-1} = 0.0127 \text{ S cm}^{-1}$ molar conductivity $\Lambda_m = \kappa \times 1000/C$ $= 0.0127 \text{ S cm}^{-1} \times 1000/0.05 \text{ mol/cm}^3$ $= 254.77 \text{ S cm}^2 \text{ mol}^{-1}$ or $A = \pi r^2$ $= 3.14 \times 0.5 \times 0.5 \text{ cm}^2$ $= 0.785 \text{ cm}^2$ $l = 45.5 \text{ cm}$ $G^* = l/A = 45.5 \text{ cm} / 0.785 \text{ cm}^2$ $= 57.96 \text{ cm}^{-1}$ $K = G^* / R$ $= 57.96 \text{ cm}^{-1} / 4.55 \times 10^3 \Omega = 1.27 \times 10^{-2} \text{ S cm}^{-1}$ $\Lambda_m = \kappa \times 1000/C$ $= [1.27 \times 10^{-2} \text{ S cm}^{-1}] \times 1000 / 0.05 \text{ mol/cm}^3$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$



	= 254.77 S cm ² mol ⁻¹	½													
12	<p>a. The particles of the dispersed phase have no affinity for the dispersion medium/solvent repelling (hating) colloidal sols. Example: metal and their sulphides</p> <p>b. The reactant and the catalyst are in the same phase.</p> $\text{CH}_3\text{COOCH}_3(\text{l}) + \text{H}_2\text{O}(\text{l}) \xrightarrow{\text{HCl}(\text{l})} \text{CH}_3\text{COOH}(\text{aq}) + \text{CH}_3\text{OH}(\text{aq})$ <p>c. oil is dispersed in water/Oil is dispersed phase and water is dispersion medium. Ex- milk</p>	½ + ½ ½ + ½ ½ + ½													
	(or any other correct example)														
	OR														
12	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Physisorption</th> <th style="text-align: center;">Chemisorption</th> <th rowspan="4" style="vertical-align: middle; text-align: center;">(1+1+1)</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">1</td> <td>Because of van der Waals forces</td> <td>Caused by chemical bond formation</td> </tr> <tr> <td style="text-align: center;">2</td> <td>Reversible</td> <td>Irreversible</td> </tr> <tr> <td style="text-align: center;">3</td> <td>Enthalpy of adsorption is low(20-40 kJ/mol)</td> <td>Enthalpy of adsorption is high(80-240)kJ/mol</td> </tr> </tbody> </table>		Physisorption	Chemisorption	(1+1+1)	1	Because of van der Waals forces	Caused by chemical bond formation	2	Reversible	Irreversible	3	Enthalpy of adsorption is low(20-40 kJ/mol)	Enthalpy of adsorption is high(80-240)kJ/mol	(Or any other correct difference)
	Physisorption	Chemisorption	(1+1+1)												
1	Because of van der Waals forces	Caused by chemical bond formation													
2	Reversible	Irreversible													
3	Enthalpy of adsorption is low(20-40 kJ/mol)	Enthalpy of adsorption is high(80-240)kJ/mol													
13	<p>Given : T_b of glucose solution= 100.20°C</p> $\Delta T_b = K_b \cdot m$ $m = 0.20 / 0.512$ $m = 0.390 \text{ mol/kg}$ $\Delta T_f = K_f \cdot m$ $\Delta T_f = 1.86 \text{ K kg/mol} \times 0.390 \text{ mol/kg}$ $\Delta T_f = 0.725 \text{ K}$ <p>Freezing point of solution = 273.15K – 0.725 = 272.425K</p>	1 ½ ½ 1													
14	<p>a. Metal is converted into a volatile compound which on strong heating decomposes to give pure metal.</p> <p>b. It selectively prevents one of the sulphide ores from coming to the froth.</p> <p>c. Coke</p>	1 1 1													
15	<p>a. For bcc structure</p> $a = 4r / \sqrt{3} \quad \text{or} \quad r = \sqrt{3}a/4$ $r = \sqrt{3} \times 400 \text{ pm} / 4$	½													

	= $1.732 \times 400 \text{ pm}/4$ = 173.2 pm b. (i) Impurity defect (ii) Cationic vacancies are created.	$\frac{1}{2}$ 1 1
16	a. Due to steric hindrance and +I effect caused by two alkyl groups in propanone. b. Due to electron withdrawing nature of $-\text{NO}_2$ group which increases the acidic strength and decreases the pK_a value . c. $(\text{CH}_3)_2\text{CH-CHO}$ has one α -H atom whereas α - H atom is absent in $(\text{CH}_3)_3\text{C-CHO}$.	$\frac{1}{2} + \frac{1}{2}$ 1 1
17	a. Ethylene Glycol and Terephthalic acid $\text{HOH}_2\text{C-CH}_2\text{OH}$, p- $\text{HOOC-C}_6\text{H}_4\text{-COOH}$ b. Tetrafluoroethene , $\text{CF}_2=\text{CF}_2$ c. Hexamethylenediamine and adipic acid $\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$, $\text{HOOC}(\text{CH}_2)_4\text{ COOH}$	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
18	a. It is the magnitude of difference in energy between the two sets of d orbital i.e. t_{2g} and e_g $t_{2g}^3 e_g^1$ b. In $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$, $\text{Ni}^{+2}(3d^8)$ has two unpaired electrons which do not pair up in the presence of weak field ligand H_2O .	1 1 1
19	a. $(\text{CH}_3)_3\text{C-OH}$ undergoes dehydration.  b. Methyl group is introduced at ortho and para positions.  c. Phenol is converted to benzene. 	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$

20	a.  b.  c. 	1,1,1
21	a. In CuCl_2 , Cu is in +2 oxidation state which is more stable due to high hydration enthalpy as compared to Cu_2Cl_2 in which Cu is in +1 oxidation state b. Due to lanthanoid contraction c. Because HCl is oxidised to chlorine.	1 1 1
22	a. Neurologically active drugs / chemical compounds used for treatment of stress / anxiety and mild or even severe mental diseases. b. Anionic detergents are sodium salts of sulphonated long chain alcohols or hydrocarbons / alkylbenzene sulphonate or detergents whose anionic part is involved in cleansing action. c. Disinfectants kill or prevent growth of microbes and are applied on inanimate / non living objects	1 1 1
23	(i)Concerned , caring, socially alert, leadership (or any other 2 values) (ii)starch (iii) α -Helix and β -pleated sheets (iv)Vitamin B / B ₁ / B ₂ / B ₆ / C (any two)	$\frac{1}{2} + \frac{1}{2}$ 1 $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
24	$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{40} \log \frac{100}{25}$ $= \frac{2.303}{40} \log 4$ $= \frac{2.303}{40} \times 0.6021$ $k = 0.0347 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{k}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

	$t_{1/2} = \frac{0.693}{0.0347 \text{ min}^{-1}} = 19.98 \text{ min} = 20 \text{ min}$ b. (i) First order reaction (ii) Zero order reaction OR	1 1 1
24	(a) $\text{Rate} = k [\text{NO}]^x [\text{O}_2]^y$ $7.2 \times 10^{-2} = k[0.3]^x [0.2]^y \text{ ----- Eqn (1)}$ $6.0 \times 10^{-3} = k[0.1]^x [0.1]^y \text{ ----- Eqn (2)}$ $2.88 \times 10^{-1} = k[0.3]^x [0.4]^y \text{ ----- Eqn (3)}$ $2.40 \times 10^{-2} = k[0.4]^x [0.1]^y \text{ ----- Eqn (4)}$ Dividing eqn 4 by eqn 2 $\frac{2.40 \times 10^{-2}}{6.0 \times 10^{-3}} = \frac{k[0.4]^x [0.1]^y}{k[0.1]^x [0.1]^y}$ $x=1$ Dividing eqn 3 by eqn 1 $\frac{2.88 \times 10^{-1}}{7.2 \times 10^{-2}} = \frac{k[0.3]^x [0.4]^y}{k[0.3]^x [0.2]^y}$ $y = 2$ order w.r.t. NO = 1, order w.r.t O ₂ is 2 (b) Rate law $\text{Rate} = k [\text{NO}]^1 [\text{O}_2]^2$, over all order of the reaction is 3. c. Rate constant $k = \frac{\text{rate}}{[\text{NO}]^1 [\text{O}_2]^2} = \frac{7.2 \times 10^{-2}}{0.3 \times (0.2)^2}$ $k = 6.0 \text{ mol}^{-2} \text{ L}^2 \text{ min}^{-1}$	1 1 ½ , ½ ½ + ½ 1
25	a. (i) Thermal stability of hydrides decreases down the group/ Bond dissociation enthalpy decreases down the group. (ii) Because Cl ₂ in presence of moisture liberates nascent oxygen. (iii) Interatomic interactions are weak b.(i) (ii) 	1 1 1 1,1
	OR	




25	a) Size of Nitrogen is smaller than Chlorine. b) $2\text{F}_2 + 2\text{H}_2\text{O} \rightarrow 4\text{HF} + \text{O}_2$ / HF and O ₂ are produced c) PH ₃ / Phosphine d) XeF ₂ e) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+} + \text{NO} \rightarrow [\text{Fe}(\text{H}_2\text{O})_5(\text{NO})]^{2+} + \text{H}_2\text{O}$	1 1 1 1 1
26.	<p>The table contains five chemical structures labeled (A) through (E):</p> <ul style="list-style-type: none"> (A) Anisole: A benzene ring with an -NH₂ group attached. (B) N-methylbenzylamine: A benzene ring with an -NHCOCH₃ group attached. (C) N,N-dimethylbenzylamine: A benzene ring with an -HN(COCH₃)₂ group attached. (D) 4-nitroaniline: A benzene ring with an -NH₂ group at position 1 and an -NO₂ group at position 4. (E) Phenylammonium sulfate: A benzene ring with a phenyl group attached to an ammonium ion (NH₃⁺) and a sulfate anion (SO₄⁻). 	1x5=5
OR		

26	<p>a. i)</p>  <p>ii)</p>  <p>iii)</p> 	<p>1,1,1</p> <p>1</p> <p>1</p>
	<p>b. $C_6H_5NH_2 < C_6H_5CH_2NH_2 < CH_3NH_2 < (CH_3)_2NH$</p> <p>c. Add $NaNO_2 + HCl$ to both the compounds at 273K followed by addition of phenol. Aniline gives orange dye (or any other correct test)</p>	

1	Dr. (Mrs.) Sangeeta Bhatia	12	Sh. S. Vallabhan	
2	Dr. K.N. Uppadhyay	13	Dr. Bhagyabati Nayak	
3	Prof. R.D. Shukla	14	Ms. Anila Mechur Jayachandran	
4	Sh. S.K. Munjal	15	Mrs. Deepika Arora	
5	Sh. D.A. Mishra	16	Ms. Seema Bhatnagar	
6	Sh. Rakesh Dhawan	17	Mrs. Sushma Sachdeva	
7	Dr. (Mrs.) Sunita Ramrakhiani	18	Dr. Azhar Aslam Khan	
8	Mrs. Preeti Kiran	19	Mr. Roop Narain Chauhan	
9	Ms. Neeru Sofat	20	Mr. Mukesh Kumar Kaushik	
10	Sh. Pawan Singh Meena	21	Ms. Abha Chaudhary	
11	Mrs. P. Nirupama Shankar	22	Ms. Garima Bhutani	




Marking scheme – 2017**CHEMISTRY (043)/ CLASS XII****FOREIGN 2017 - Set - 56/2/2**








Q.NO .	VALUE POINTS	MARK S
1	2-Methylbut-3-en-2-ol	1
2	Neopentane , $\text{C}(\text{CH}_3)_4$	1
3	$\text{H}_2\text{Te} > \text{H}_2\text{Se} > \text{H}_2\text{S} > \text{H}_2\text{O}$	1
4	P_3Q_2	1
5	To make the surface available again for more reaction to occur / To remove the product formed from the surface of the catalyst.	1
6	a. Pentaamminesulphatocobalt(III) chloride b. $[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{NO}_2)]$	1 1
7	a. Zinc to silver b. Concentration of Zn^{2+} ions will increase and Ag^+ ions will decrease.	1 1
8	a. Cr^{3+} b. Mn^{3+} c. Ti^{4+} d. Mn^{3+}	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
9	a. $\text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow[\text{H}^+]{\text{H}_2\text{O}} \text{CH}_3\text{CH}(\text{OH})\text{CH}_3 \xrightarrow{\text{CrO}_3} \text{CH}_3\text{COCH}_3$ b. $\text{CH}_3\text{CH}_2\text{COOH} \xrightarrow{\text{Br}_2/\text{Red P}} \text{CH}_3\text{CH}(\text{Br})\text{COOH} \xrightarrow[\text{ii)}\text{H}^+]{\text{i)} \text{aq KOH or NaOH}} \text{CH}_3\text{CH}(\text{OH})\text{COOH}$ (or any other suitable method)	1 1
	OR	
9	a.Etard reaction: $\begin{array}{ccc} \text{Toluene} & + \text{CrO}_2\text{Cl}_2 & \xrightarrow{\text{CS}_2} \\ \text{Benzyl alcohol} & & \text{Chromium complex} \\ & & \xrightarrow{\text{H}_3\text{O}^+} \text{Benzaldehyde} \end{array}$	1



	<p style="text-align: center;">or</p> <p style="text-align: center;">  (i) $\text{CrO}_2\text{Cl}_2, \text{CS}_2$ (ii) H_3O^+ </p> <p>b.Wolff-Kishner reduction:</p> <p style="text-align: center;"> $\text{C=O} \xrightarrow[-\text{H}_2\text{O}]{\text{NH}_2\text{NH}_2} \text{C}=\text{NNH}_2 \xrightarrow[\text{heat}]{\text{KOH/ethylene glycol}} \text{CH}_2 + \text{N}_2$ </p> <p style="text-align: center;">or</p> <p style="text-align: center;"> $\text{C=O} \xrightarrow{\text{(i) NH}_2\text{NH}_2} \text{C}=\text{NNH}_2 \xrightarrow{\text{(ii) KOH/ethylene glycol, heat}} \text{CH}_2 + \text{N}_2$ </p>	
10	<p>The relative lowering of vapour pressure of a solution is equal to the mole fraction of the solute. /</p> <p>The vapour pressure of a solution of a non-volatile solute is equal to the vapour pressure of the pure solvent at that temperature multiplied by its mole fraction.</p> <p>Negative deviation due to formation of Hydrogen bond between chloroform and acetone.</p>	<p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
11	<p>a. Phenol & Formaldehyde</p> <p style="text-align: center;">  & HCHO </p> <p>b.Vinyl chloride , $\text{CH}_2=\text{CHCl}$</p> <p>c. 1,3-Butadiene & styrene</p> <p style="text-align: center;">  </p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
12	<p>a. It is the magnitude of difference in energy between the two sets of d orbital i.e. t_{2g} and e_g</p> <p>$t_{2g}^4 e_g^0$</p> <p>b. In $[\text{Ni}(\text{CN})_4]^{2-}$, CN^- is a strong field ligand and pairing takes place whereas in $[\text{NiCl}_4]^{2-}$, due to the presence of Cl^- , a weak field ligand no pairing occurs / diagrammatic representation</p>	<p>1</p> <p>1</p> <p>1</p>

13.	<p>a. $(CH_3)_3C-OH$ undergoes dehydration.</p> $\begin{array}{c} CH_3 \\ \\ CH_3-C-OH \\ \\ CH_3 \end{array} \xrightarrow[573K]{Cu} \begin{array}{c} CH_3 \\ \\ CH_3-C=CH_2 \end{array}$ <p>b. Methyl group is introduced at ortho and para positions.</p> $\begin{array}{c} OCH_3 \\ \\ C_6H_5 \end{array} + CH_3Cl \xrightarrow[CS_2]{\text{Anhyd. AlCl}_3} \begin{array}{c} OCH_3 \\ \\ C_6H_4-CH_3 \end{array} + \begin{array}{c} OCH_3 \\ \\ C_6H_4-CH_3 \end{array}$ <p>c. Phenol is converted to benzene.</p> $\begin{array}{c} OH \\ \\ C_6H_5 \end{array} + Zn \longrightarrow \begin{array}{c} \\ \\ C_6H_5 \end{array} + ZnO$	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
14	<p>a. $Eu^{2+}(4f^7)$ is a strong reducing agent because Eu^{3+} is more stable than Eu^{2+}.</p> <p>b. Dichromate ion changes to chromate ion /</p> OH^- $Cr_2O_7^{2-} \text{ (orange)} \rightarrow CrO_4^{2-} \text{ (yellow)}$ <p>c. Due to the irregular variation in ionisation enthalpies (sum of 1st and 2nd ionisation enthalpies), heat of sublimation and enthalpy of hydration/ due to irregular electronic configurations from left to right in a period which changes the ionisation potential.</p>	1 1 1
15	<p>a. Antiseptics are the chemicals which either kill or prevent growth of microbes on living tissues.</p> <p>b. Cationic detergents are quaternary ammonium salts of amines with acetates, chlorides or bromides as anions / detergents whose cationic part is involved in cleansing action.</p> <p>c. Antibiotics which kill or inhibit a wide range of Gram-positive and Gram-negative bacteria.</p>	1 1 1
16	$\begin{aligned} A &= \pi r^2 \\ &= 3.14 \times 0.5 \times 0.5 \text{ cm}^2 \\ &= 0.785 \text{ cm}^2 \\ l &= 45.5 \text{ cm} \\ \rho &= R \times A / l \\ \rho &= 4.55 \times 10^3 \Omega \times 0.785 \text{ cm}^2 / 45.5 \text{ cm} \\ \rho &= 78.5 \Omega \text{ cm} \end{aligned}$ <p>conductivity, $\kappa = 1/\rho$</p> $\kappa = 1/78.5 \text{ S cm}^{-1} = 0.0127 \text{ S cm}^{-1}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

	$\text{molar conductivity } \Lambda_m = \kappa \times 1000/C$ $= 0.0127 \text{ S cm}^{-1} \times 1000/0.05 \text{ mol/cm}^3$ $= 254.77 \text{ S cm}^2 \text{ mol}^{-1}$ or $A = \pi r^2$ $= 3.14 \times 0.5 \times 0.5 \text{ cm}^2$ $= 0.785 \text{ cm}^2$ $l = 45.5 \text{ cm}$ $G^* = l/A = 45.5 \text{ cm} / 0.785 \text{ cm}^2$ $= 57.96 \text{ cm}^{-1}$ $K = G^* / R$ $= 57.96 \text{ cm}^{-1} / 4.55 \times 10^3 \Omega = 1.27 \times 10^{-2} \text{ S cm}^{-1}$ $\Lambda_m = \kappa \times 1000/C$ $= [1.27 \times 10^{-2} \text{ S cm}^{-1}] \times 1000 / 0.05 \text{ mol/cm}^3$ $= 254.77 \text{ S cm}^2 \text{ mol}^{-1}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$												
17	a. The particles of the dispersed phase have no affinity for the dispersion medium/solvent repelling (hating) colloidal sols. Example: metal and their sulphides b. The reactant and the catalyst are in the same phase. $\text{CH}_3\text{COOCH}_3(\text{l}) + \text{H}_2\text{O}(\text{l}) \xrightarrow{\text{HCl}(\text{l})} \text{CH}_3\text{COOH}(\text{aq}) + \text{CH}_3\text{OH}(\text{aq})$ c. Oil is dispersed in water/Oil is dispersed phase and water is dispersion medium. Ex- milk (or any other correct example)	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$												
	OR													
17	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Physisorption</th> <th style="text-align: center;">Chemisorption</th> </tr> </thead> <tbody> <tr> <td>1</td><td>Because of van der Waals forces</td><td>Caused by chemical bond formation</td></tr> <tr> <td>2</td><td>Reversible</td><td>Irreversible</td></tr> <tr> <td>3</td><td>Enthalpy of adsorption is low(20-40 kJ/mol)</td><td>Enthalpy of adsorption is high(80-240)kJ/mol</td></tr> </tbody> </table> (Or any other correct difference)		Physisorption	Chemisorption	1	Because of van der Waals forces	Caused by chemical bond formation	2	Reversible	Irreversible	3	Enthalpy of adsorption is low(20-40 kJ/mol)	Enthalpy of adsorption is high(80-240)kJ/mol	$1+1+1$
	Physisorption	Chemisorption												
1	Because of van der Waals forces	Caused by chemical bond formation												
2	Reversible	Irreversible												
3	Enthalpy of adsorption is low(20-40 kJ/mol)	Enthalpy of adsorption is high(80-240)kJ/mol												
18	Given : T_b of glucose solution= 100.20°C $\Delta T_b = K_b \cdot m$													

	$m = 0.20 / 0.512$ $m = 0.390 \text{ mol/kg}$ $\Delta T_f = K_f \cdot m$ $\Delta T_f = 1.86 \text{ K kg/mol} \times 0.390 \text{ mol/kg}$ $\Delta T_f = 0.725 \text{ K}$ Freezing point of solution = $273.15 \text{ K} - 0.725$ $= 272.425 \text{ K}$	1 ½ ½ 1
19	a) Zone Refining – Impurities are more soluble in the melt than in the solid metal. b) Collectors enhance non-wettability of the mineral particles.Ex Pine oil/ fatty acids c) Carbon monoxide (CO)	1 1 1
20	a. For bcc structure $a = 4r / \sqrt{3}$ or $r = \sqrt{3}a/4$ $r = \sqrt{3} \times 400 \text{ pm} / 4$ $= 1.732 \times 400 \text{ pm}/4$ $= 173.2 \text{ pm}$ b. (i) Impurity defect (ii) Cationic vacancies are created.	½ ½ 1 1
21	a.  b.  c. 	1,1,1
22	a. Due to steric hindrance and +I effect caused by two alkyl groups in propanone. b. Due to electron withdrawing nature of $-NO_2$ group which increases the acidic strength and decreases the pK_a value . c. $(CH_3)_2CH-CHO$ has one α -H atom whereas α - H atom is absent in $(CH_3)_3C-CHO$.	½+ ½ 1 1
23	(i)Concerned , caring, socially alert, leadership (or any other 2 values) (ii)starch	½ + ½ 1

	(iii) α -Helix and β -pleated sheets (iv) Vitamin B / B ₁ / B ₂ / B ₆ / C (any two)	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
24	a. (i) Thermal stability of hydrides decreases down the group/ Bond dissociation enthalpy decreases down the group. (ii) Because Cl ₂ in presence of moisture liberates nascent oxygen. (iii) Interatomic interactions are weak b.(i) (ii)	1 1 1  
	OR	
24	a) Size of nitrogen is smaller than Chlorine. b) $2\text{F}_2 + 2\text{H}_2\text{O} \rightarrow 4\text{HF} + \text{O}_2$ / HF and O ₂ are produced c) PH ₃ /Phosphine d) XeF ₂ e) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+} + \text{NO} \rightarrow [\text{Fe}(\text{H}_2\text{O})_5(\text{NO})]^{2+} + \text{H}_2\text{O}$	1 1 1 1 1
25	(A)  (B)  (C)  (D)  (E) 	1×5=5
	OR	


25	<p>a. i)</p>  <p>ii)</p>  <p>iii)</p> 	1,1,1
	<p>b. $C_6H_5NH_2 < C_6H_5CH_2NH_2 < CH_3NH_2 < (CH_3)_2NH$</p> <p>c. Add $NaNO_2 + HCl$ to both the compounds at 273K followed by addition of phenol. Aniline gives orange dye (or any other correct test)</p>	1 1
26.	$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$ $= \frac{2.303}{40} \log \frac{100}{25}$ $= \frac{2.303}{40} \log 4$ $= \frac{2.303}{40} \times 0.6021$ $k = 0.0347 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{k}$ $t_{1/2} = \frac{0.693}{0.0347 \text{ min}^{-1}} = 19.98 \text{ min} = 20 \text{ min}$ <p>b. (i) first order reaction (ii) zero order reaction</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
	OR	
26	<p>(a)</p> $\text{Rate} = k [NO]^x [O_2]^y$ $7.2 \times 10^{-2} = k[0.3]^x [0.2]^y \text{ ----- Eqn (1)}$ $6.0 \times 10^{-3} = k[0.1]^x [0.1]^y \text{ ----- Eqn (2)}$ $2.88 \times 10^{-1} = k[0.3]^x [0.4]^y \text{ ----- Eqn (3)}$ $2.40 \times 10^{-2} = k[0.4]^x [0.1]^y \text{ ----- Eqn (4)}$	

<p>Dividing eqn 4 by eqn 2</p> $\frac{2.40 \times 10^{-2}}{6.0 \times 10^{-3}} = \frac{k[0.4]^x [0.1]^y}{k[0.1]^x [0.1]^y}$ $4 = \frac{[0.4]^x}{[0.1]^x} = 4^x$ $4 = 4^x \Rightarrow x=1$ <p>Dividing eqn 3 by eqn 1</p> $\frac{2.88 \times 10^{-1}}{7.2 \times 10^{-2}} = \frac{k[0.3]^x [0.4]^y}{k[0.3]^x [0.2]^y}$ $4 = \frac{[0.4]^y}{[0.2]^y} = 4^y$ $4 = 2^y \Rightarrow y = 2$ <p>order w.r.t. NO = 1, order w.r.t O₂ is 2</p> <p>(b) Rate law</p> <p>Rate = k [NO]¹ [O₂]² ;The overall order of the reaction is 3.</p> <p>c. rate constant k = $\frac{\text{rate}}{[\text{NO}]^1 [\text{O}_2]^2} = \frac{7.2 \times 10^{-2}}{0.3 \times (0.2)^2}$ $k = 6.0 \text{ mol}^{-2} \text{ L}^2 \text{ min}^{-1}$</p>	<p>1</p> <p>1</p> <p>½ , ½</p> <p>½ + ½</p> <p>1</p>
--	--




1	Dr. (Mrs.) Sangeeta Bhatia	12	Sh. S. Vallabhan	
2	Dr. K.N. Uppadhyा	13	Dr. Bhagyabati Nayak	
3	Prof. R.D. Shukla	14	Ms. Anila Mechur Jayachandran	
4	Sh. S.K. Munjal	15	Mrs. Deepika Arora	
5	Sh. D.A. Mishra	16	Ms. Seema Bhatnagar	
6	Sh. Rakesh Dhawan	17	Mrs. Sushma Sachdeva	
7	Dr. (Mrs.) Sunita Ramrakhiani	18	Dr. Azhar Aslam Khan	
8	Mrs. Preeti Kiran	19	Mr. Roop Narain Chauhan	
9	Ms. Neeru Sofat	20	Mr. Mukesh Kumar Kaushik	
10	Sh. Pawan Singh Meena	21	Ms. Abha Chaudhary	
11	Mrs. P. Nirupama Shankar	22	Ms. Garima Bhutani	




Marking scheme – 2017**CHEMISTRY (043)/ CLASS XII****FOREIGN 2017 - Set - 56/2/3**



Q.NO .	VALUE POINTS	MARK S
1	$\text{H}_2\text{Te} > \text{H}_2\text{Se} > \text{H}_2\text{S} > \text{H}_2\text{O}$	1
2	To make the surface available again for more reaction to occur / To remove the product formed from the surface of the catalyst.	1
3	2-Phenylpropan-2-ol	1
4	Neopentane , $\text{C}(\text{CH}_3)_4$	1
5	P_3Q_2	1
6	a. Zinc to silver b. Concentration of Zn^{2+} ions will increase and Ag^+ ions will decrease.	1 $\frac{1}{2} + \frac{1}{2}$
7	a. Cr^{3+} b. Mn^{3+} c. Ti^{4+} d. Mn^{3+}	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
8	a. $\text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow[\text{H}^+]{\text{H}_2\text{O}} \text{CH}_3\text{CH}(\text{OH})\text{CH}_3 \xrightarrow{\text{CrO}_3} \text{CH}_3\text{COCH}_3$ b. $\text{CH}_3\text{CH}_2\text{COOH} \xrightarrow{\text{Br}_2/\text{Red P}} \text{CH}_3\text{CH}(\text{Br})\text{COOH} \xrightarrow[\text{ii) H}^+]{\text{i) aq KOH or NaOH}} \text{CH}_3\text{CH}(\text{OH})\text{COOH}$ (or any other suitable method)	1 1
	OR	
8	a.Etard reaction: $\text{Toluene} + \text{CrO}_2\text{Cl}_2 \xrightarrow{\text{CS}_2} \text{Chromium complex} \xrightarrow{\text{H}_3\text{O}^+} \text{Benzaldehyde}$ or	1

	<p style="text-align: center;">  b. Wolff-Kishner reduction: $\text{C=O} \xrightarrow[-\text{H}_2\text{O}]{\text{NH}_2\text{NH}_2} \text{C=NNH}_2 \xrightarrow[\text{heat}]{\text{KOH/ethylene glycol}} \text{CH}_2 + \text{N}_2$ or $\text{C=O} \xrightarrow{\text{(i) NH}_2\text{NH}_2, \text{(ii) KOH/ethylene glycol, heat}} \text{CH}_2 + \text{N}_2$ </p>	1
9	The increase in boiling point of the solvent in a solution when a non-volatile solute is added. Because it depends upon molality / the number of solute particles rather than their nature/ $\Delta T_b \propto m$	1 1
10	a. Tetraamminechloridonitrito-N-cobalt(III) chloride b. $[\text{CoCl}_2(\text{en})_2]\text{Cl}$	1 1
11	a. In CuCl_2 , Cu is in +2 oxidation state which is more stable due to high hydration enthalpy as compared to Cu_2Cl_2 in which Cu is in +1 oxidation state b. Due to lanthanoid contraction c. Because HCl is oxidised to chlorine.	1 1 1
12	a. Drugs that reduce or abolish pain without causing impairment of consciousness , mental confusion or paralysis. b. Anionic detergents are sodium salts of sulphonated long chain alcohols or hydrocarbons / alkylbenzene sulphonate or detergents whose anionic part is involved in cleansing action. c. Antacids are chemical compounds which are used for the treatment of excess acid produced in the stomach.	1 1 1
13	$\begin{aligned} A &= \pi r^2 \\ &= 3.14 \times 0.5 \times 0.5 \text{ cm}^2 \\ &= 0.785 \text{ cm}^2 \end{aligned}$ $\begin{aligned} l &= 45.5 \text{ cm} \\ \rho &= R \times A / l \\ \rho &= 4.55 \times 10^3 \Omega \times 0.785 \text{ cm}^2 / 45.5 \text{ cm} \\ \rho &= 78.5 \Omega \text{ cm} \end{aligned}$ conductivity , $\kappa = 1 / \rho$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

	$= 1/78.5 \text{ S cm}^{-1} = 0.0127 \text{ S cm}^{-1}$ molar conductivity $\Lambda_m = \kappa \times 1000/C$ $= 0.0127 \text{ S cm}^{-1} \times 1000/0.05 \text{ mol/cm}^3$ $= 254.77 \text{ S cm}^2 \text{ mol}^{-1}$ or $A = \pi r^2$ $= 3.14 \times 0.5 \times 0.5 \text{ cm}^2$ $= 0.785 \text{ cm}^2$ $l = 45.5 \text{ cm}$ $G^* = l/A = 45.5 \text{ cm} / 0.785 \text{ cm}^2$ $= 57.96 \text{ cm}^{-1}$ $K = G^*/R$ $= 57.96 \text{ cm}^{-1} / 4.55 \times 10^3 \Omega = 1.27 \times 10^{-2} \text{ S cm}^{-1}$ $\Lambda_m = \kappa \times 1000/C$ $= [1.27 \times 10^{-2} \text{ S cm}^{-1}] \times 1000 / 0.05 \text{ mol/cm}^3$ $= 254.77 \text{ S cm}^2 \text{ mol}^{-1}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$												
14	<p>a. The particles of the dispersed phase have no affinity for the dispersion medium/solvent repelling (hating) colloidal sols. Example: metal and their sulphides</p> <p>b. The reactant and the catalyst are in the same phase.</p> $\text{CH}_3\text{COOCH}_3(\text{l}) + \text{H}_2\text{O}(\text{l}) \xrightarrow{\text{HCl}(\text{l})} \text{CH}_3\text{COOH}(\text{aq}) + \text{CH}_3\text{OH}(\text{aq})$ <p>c. oil is dispersed in water/Oil is dispersed phase and water is dispersion medium. Ex- milk (or any other correct example)</p>	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$												
	OR													
14	<table border="1"> <thead> <tr> <th></th> <th>Physisorption</th> <th>Chemisorption</th> </tr> </thead> <tbody> <tr> <td>1</td><td>Because of van der Waals forces</td><td>Caused by chemical bond formation</td></tr> <tr> <td>2</td><td>Reversible</td><td>Irreversible</td></tr> <tr> <td>3</td><td>Enthalpy of adsorption is low(20-40 kJ/mol)</td><td>Enthalpy of adsorption is high(80-240)kJ/mol</td></tr> </tbody> </table> <p style="text-align: center;">(Or any other correct difference)</p>		Physisorption	Chemisorption	1	Because of van der Waals forces	Caused by chemical bond formation	2	Reversible	Irreversible	3	Enthalpy of adsorption is low(20-40 kJ/mol)	Enthalpy of adsorption is high(80-240)kJ/mol	$1+1+1$
	Physisorption	Chemisorption												
1	Because of van der Waals forces	Caused by chemical bond formation												
2	Reversible	Irreversible												
3	Enthalpy of adsorption is low(20-40 kJ/mol)	Enthalpy of adsorption is high(80-240)kJ/mol												




15	a.  b.  c. 	1,1,1
16	Given : T_b of glucose solution = 100.20°C $\Delta T_b = K_b \cdot m$ $m = 0.20 / 0.512$ $m = 0.390 \text{ mol/kg}$ $\Delta T_f = K_f \cdot m$ $\Delta T_f = 1.86 \text{ K kg/mol} \times 0.390 \text{ mol/kg}$ $\Delta T_f = 0.725 \text{ K}$ Freezing point of solution = $273.15 \text{ K} - 0.725$ = 272.425 K	1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
17	a.(i) Vapour phase refining/ van Arkel method (ii) Zone refining (iii) Electrolytic refining b.(i) Froth floatation process (ii) Magnetic separation (iii) Leaching	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
18	a. For bcc structure $a = 4r / \sqrt{3}$ or $r = \sqrt{3}a/4$ $r = \sqrt{3} \times 400 \text{ pm} / 4$ = $1.732 \times 400 \text{ pm}/4$ = 173.2 pm b. (i) Impurity defect (ii) Cationic vacancies are created.	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
19	a. Due to steric hindrance and +I effect caused by two alkyl groups in propanone. b. Due to electron withdrawing nature of $-\text{NO}_2$ group which increases the acidic strength and decreases the pK_a value . c. $(\text{CH}_3)_2\text{CH-CHO}$ has one α -H atom whereas α - H atom is absent in $(\text{CH}_3)_3\text{C-CHO}$.	$\frac{1}{2} + \frac{1}{2}$ 1 1
20	a. Chloroprene , $\text{CH}_2=\text{C}(\text{Cl})-\text{CH}=\text{CH}_2$ b. 1,3- Butadiene & Acrylonitrile $\text{CH}_2=\text{CH-CH=CH}_2$ & $\text{CH}_2=\text{CHCN}$	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$

	c. 3-Hydroxybutanoic acid & 3-Hydroxypentanoic acid $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{COOH}$ & $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{COOH}$	$\frac{1}{2} + \frac{1}{2}$
21	a) It is the magnitude of difference in energy between the two sets of d orbital i.e. t_{2g} and e_g $t^4_{2g} e_g^0$ b) sp^3d^2 , paramagnetic	1 1 $\frac{1}{2} + \frac{1}{2}$
22	a. Methanol and 2-methyl-2-iodopropane are formed. $\begin{array}{ccc} \text{CH}_3 & & \text{CH}_3 \\ & & \\ \text{CH}_3-\text{C}-\text{O}-\text{CH}_3 & + \text{HI} \longrightarrow & \text{CH}_3\text{OH} + \text{CH}_3-\text{C}-\text{I} \\ & & \\ \text{CH}_3 & & \text{CH}_3 \end{array}$ b. 2-Methoxy acetophenone and 4-Methoxy acetophenone are formed $\begin{array}{ccc} \text{OCH}_3 & & \text{OCH}_3 \\ & & \\ \text{C}_6\text{H}_5 & + \text{CH}_3\text{COCl} \xrightarrow{\text{Anhyd. AlCl}_3} & \text{C}_6\text{H}_5\text{COCH}_3 + \text{C}_6\text{H}_5\text{COCH}_3 \\ & & \\ \text{OCH}_3 & & \text{OCH}_3 \end{array}$ c. o-Bromophenol and p-Bromophenol are formed. $\begin{array}{ccc} \text{OH} & & \text{OH} \\ & & \\ \text{C}_6\text{H}_5 & \xrightarrow[\text{273 K}]{\text{Br}_2 \text{ in } \text{CS}_2} & \text{C}_6\text{H}_5\text{Br} + \text{C}_6\text{H}_5\text{Br} \\ & & \\ \text{OH} & & \text{Br} \end{array}$	1 1 1
	(Award full marks if the student writes only equation)	
23	(i)Concerned , caring, socially alert, leadership (or any other 2 values) (ii)starch (iii) α -Helix and β -pleated sheets (iv)Vitamin B / B ₁ / B ₂ / B ₆ / C (any two)	$\frac{1}{2} + \frac{1}{2}$ 1 $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
24	(A)  (B)  (C) 	1×5=5



OR

24



1,1,1



1

c. Add $NaNO_2 + HCl$ to both the compounds at 273K followed by addition of phenol. Aniline gives orange dye
(or any other correct test)

1

25.

$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$$

 $\frac{1}{2}$

$$= \frac{2.303}{40} \log \frac{100}{25}$$

 $\frac{1}{2}$

$$= \frac{2.303}{40} \log 4$$

$$= \frac{2.303}{40} \times 0.6021$$



$$k = 0.0347 \text{ min}^{-1}$$

 $\frac{1}{2}$

$$t_{1/2} = \frac{0.693}{k}$$

 $\frac{1}{2}$

	$t_{1/2} = \frac{0.693}{0.0347 \text{ min}^{-1}} = 19.98 \text{ min} = 20 \text{ min}$ b. (i) first order reaction (ii) zero order reaction	1 1 1
	OR	
25	(a) $\text{Rate} = k [NO]^x [O_2]^y$ $7.2 \times 10^{-2} = k[0.3]^x [0.2]^y \text{ ----- Eqn (1)}$ $6.0 \times 10^{-3} = k[0.1]^x [0.1]^y \text{ ----- Eqn (2)}$ $2.88 \times 10^{-1} = k[0.3]^x [0.4]^y \text{ ----- Eqn (3)}$ $2.40 \times 10^{-2} = k [0.4]^x [0.1]^y \text{ ----- Eqn (4)}$ Dividing eqn 4 by eqn 2 $\frac{2.40 \times 10^{-2}}{6.0 \times 10^{-3}} = \frac{k[0.4]^x [0.1]^y}{k[0.1]^x [0.1]^y}$ $4 = 4^x$ $x=1$ Dividing eqn 3 by eqn 1 $\frac{2.88 \times 10^{-1}}{7.2 \times 10^{-2}} = \frac{k[0.3]^x [0.4]^y}{k[0.3]^x [0.2]^y}$ $4 = 2^y$ $y=2$ order w.r.t. NO = 1, order w.r.t O ₂ is 2	1 1 1/2 , 1/2
	(b) Rate law $\text{Rate} = k [NO]^1 [O_2]^2$, The overall order of the reaction is 3. c. rate constant $k = \frac{\text{rate}}{[NO]^1 [O_2]^2} = \frac{7.2 \times 10^{-2}}{0.3 \times (0.2)^2}$ $k = 6.0 \text{ mol}^{-2} \text{ L}^2 \text{ min}^{-1}$	1/2 + 1/2
26.	a. (i) Thermal stability of hydrides decreases down the group/ Bond dissociation enthalpy decreases down the group. (ii) Because Cl ₂ in presence of moisture liberates nascent oxygen. (iii) Interatomic interactions are weak	1 1 1
	b.(i) (ii)	

	 	1,1
OR		
26	a) Size of nitrogen is smaller than Chlorine. b) $2\text{F}_2 + 2\text{H}_2\text{O} \rightarrow 4\text{HF} + \text{O}_2$ / HF and O ₂ are produced c) PH ₃ / Phosphine d) XeF ₂ e) $[\text{Fe}(\text{H}_2\text{O})_6]^{2+} + \text{NO} \rightarrow [\text{Fe}(\text{H}_2\text{O})_5(\text{NO})]^{2+} + \text{H}_2\text{O}$	1 1 1 1 1

1	Dr. (Mrs.) Sangeeta Bhatia	12	Sh. S. Vallabhan	
2	Dr. K.N. Uppadhyay	13	Dr. Bhagyabati Nayak	
3	Prof. R.D. Shukla	14	Ms. Anila Mechur Jayachandran	
4	Sh. S.K. Munjal	15	Mrs. Deepika Arora	
5	Sh. D.A. Mishra	16	Ms. Seema Bhatnagar	
6	Sh. Rakesh Dhawan	17	Mrs. Sushma Sachdeva	
7	Dr. (Mrs.) Sunita Ramrakhiani	18	Dr. Azhar Aslam Khan	
8	Mrs. Preeti Kiran	19	Mr. Roop Narain Chauhan	
9	Ms. Neeru Sofat	20	Mr. Mukesh Kumar Kaushik	
10	Sh. Pawan Singh Meena	21	Ms. Abha Chaudhary	
11	Mrs. P. Nirupama Shankar	22	Ms. Garima Bhutani	

