






**MARKING SCHEME CHEMISTRY****(CODE NO. : 56/3/C)**




Q	Value points	Mark
1	White phosphorous, because of angular strain in P <sub>4</sub> molecule/ discrete tetrahedral unit.	½ , ½
2	$\begin{array}{c} \text{CH}_3 \\   \\ \text{H}_3\text{C}-\text{C}-\text{Br} \\   \\ \text{CH}_3 \end{array}$ <p>Because carbocation intermediate derived from (CH<sub>3</sub>)<sub>3</sub> CBr is more stable than carbocation from CH<sub>3</sub>CH<sub>2</sub>Br.</p>	½  ½
3	(i) Electrophoresis (ii) by mixing two oppositely charged sols (iii) by boiling (iv) by persistent dialysis (v) by addition of electrolyte (any one)	1
4	X <sub>4</sub> Y <sub>3</sub>	1
5	2-Methylpropane-1,3-diol	1
6	Greater number of unpaired electrons, greater the interatomic interactions leading to strong metallic bonding. Zn ,no unpaired electrons hence weak metallic bonding.	1  ½ ,½
7	(i) pentaamminenitrito-N-cobalt(III) nitrate (ii) K <sub>2</sub> [Ni(CN) <sub>4</sub> ]	1 1
8	(a) H <sup>+</sup> (aq) + e <sup>-</sup> → 1/2H <sub>2</sub> (g) E° = 0.00 V is feasible at cathode because its reduction potential is higher than the other reaction.  b. Because the overall reaction doesn't involve any ion in the solution whose concentration changes during its lifetime.	½ , ½  1
9	(i) CH <sub>3</sub> MgBr , H <sub>3</sub> O <sup>+</sup> (ii) Cl <sub>2</sub> , P	1 1
10	It states that solubility of gas in liquid is directly proportional to partial pressure of the gas in equilibrium with the solution.  With increase in temperature K <sub>H</sub> value increases but solubility of gas in liquid decreases. / K <sub>H</sub> ∝ 1/solubility  OR	1  1
10	It states for solution containing volatile components the partial vapor pressure of each component of the solution is directly proportional to its mole fraction present in the solution.	1




	Ideal Solution	Non Ideal	
	1. It obeys Raoult's Law over entire range of concentration of solution.	It does not obey Raoult's Law.	
	2. Solute – Solvent interaction is nearly same as in pure solvent.	Solute – Solvent interaction is not same as solute-solute or solvent –solvent interactions.	½ + ½
	(or any other correct difference)		
11	(i) van Arkel method (ii) CO acts as reducing agent (iii) Because $\Delta S$ becomes more positive, and $\Delta G$ becomes negative.		1 1 1
12	(a)(i) Because actinoids are radioactive and show wide range of oxidation states. (ii) Transition metals form complex compounds due to small size, high ionic charge, availability of d orbitals.  b. $2\text{MnO}_4^- + 6\text{H}^+ + 5\text{SO}_3^{2-} \rightarrow 5\text{SO}_4^{2-} + 3\text{H}_2\text{O} + 2\text{Mn}^{2+}$		1 1 1
13	 <p>(i)  cis  trans</p> <p>(ii) <math>t_2g^4</math> / diagram. (iii) <math>[\text{NiCl}_4]^{2-}</math> - Chloride ion being weak field ligand does not pair d electrons. while in <math>[\text{Ni}(\text{CO})_4]</math>, CO being strong field ligand pairs up the d electrons.</p>		1 1 1
14	(i) $\text{CH}_3 - \text{CH}(\text{OH}) - \text{CN}$ (ii) $\text{C}_6\text{H}_5\text{COOH}$ (iii) $\text{CH}_3\text{CONH}_2$		1 1 1
15	 <p>(i) Caprolactum</p>		1

	(ii) Phenol + Formaldehyde  + HCHO (iii) 1,3-Butadiene + Acrylonitrile $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + \text{CH}_2=\text{CH}-\text{CN}$ (Note: half mark for structure/s and half mark for name/s)	1 1
16	(i) Starch (ii) Native Protein found in a biological system with a unique 3-D structure and biological activity is called a native protein. Denatured protein is the protein with no biological activity. (iii) Vitamin-K	1 1 1
17	$\Delta T_f = i \times K_f \times m$ For $\text{CaCl}_2$ $i = 3$ $\Delta T_f = (i \times K_f \times W_B \times 1000) / (M_B \times W_A)$ $2 = 3 \times 1.86 \times W_B \times 1000 / 111 \times 500$ $W_B = 19.89 \text{ g}$	1/2 1/2 1 1
18	$d = Z \times M / a^3 \times N_o$ $10 \text{ g/cm}^3 = Z \times 81 \text{ g/mol} / (3 \times 10^{-8} \text{ cm})^3 \times (6.023 \times 10^{23} / \text{mol})$ $Z = 2.007$ Nature of cubic unit cell = bcc	1/2 1/2 1 1
19	(i) $\text{CH}_3-\text{CH}=\text{CH}_2 \xrightarrow[\text{peroxide}]{\text{HBr}} \text{CH}_3-\text{CH}_2-\text{CH}_2-\text{Br} \xrightarrow{\text{AgF}} \text{CH}_3\text{CH}_2\text{CH}_2\text{F}$ (ii) 	1 1

19	<p style="text-align: center;"> <math>\text{PCl}_3/\text{PCl}_5</math>                      <math>\text{KCN}</math>            (iii) <math>\text{C}_2\text{H}_5\text{OH} \xrightarrow{\hspace{2cm}} \text{C}_2\text{H}_5\text{Cl} \xrightarrow{\hspace{2cm}} \text{C}_2\text{H}_5\text{CN}</math> </p> <p style="text-align: center;"><b>OR</b></p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>(i) <math>\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2</math></p> <p>(iii) <math>\text{CH}_3\text{NC}</math></p> </div> <div style="text-align: center;"> <p>(ii)</p>  </div> </div>	1           1,1,1
20	<p>(i) Because <math>-\text{NO}_2</math> is an electron withdrawing group.</p> <p>(ii) Due to H-Bonding</p> <p>(iii) Reaction occurs by <math>\text{S}_{\text{N}}1</math> mechanism., <math>3^\circ</math>-carbocation <math>(\text{CH}_3)_3\text{C}^+</math> is more stable than <math>\text{CH}_3^+</math></p>	1 1 1
21	$E^\circ_{\text{cell}} = E_{\text{R}}^\circ - E_{\text{L}}^\circ$ $= 0.00 - (-0.14)$ $E^\circ_{\text{cell}} = +0.14\text{V}$ $E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.059\text{V}}{n} \log \frac{[\text{Sn}^{2+}]}{[\text{H}^+]^2}$ $E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.059\text{V}}{2} \log \frac{[0.001]}{[0.01]^2}$ $= +0.14 - 0.0295\text{V} \log 10$ $E_{\text{cell}} = 0.1105\text{V}$	1  1  1
22	<p>(i) Because physisorption is exothermic process, so it decreases with increase in temperature.</p> <p>(ii) Because alum coagulates the impurities present in water.</p> <p>(iii) Due to continuous unbalanced bombardment / zig-zag motion of particles by the molecules of dispersion medium/ it does not allow the particles to settle down.</p>	1 1 1
23	<p>(i) Concern, Compassion, caring, empathy (any two)</p> <p>(ii) By organizing rallies , posters, street play, public speech(any other relevant answer)</p> <p>(iii) Anti depressant drugs are those which inhibit depression E.g. Iproniazide , Phenelzine (or any other)</p> <p>(iv) Saccharine / Sucralose/Alitame/Aspartame(any one)</p>	$\frac{1}{2}$ , $\frac{1}{2}$ 1   $\frac{1}{2}$ , $\frac{1}{2}$ 1
24	<p>a. )Rate of reaction is defined as change in concentration of reactants or products per unit time. Factors: concentration of reactant, temperature, pressure, surface area (any two)</p>	1  $\frac{1}{2}$ , $\frac{1}{2}$

24	<p>b. <math>\log(k_2/k_1) = E_a/2.303 R [1/T_1 - 1/T_2]</math></p> <p><math>\log(8 \times 10^{-2}/4 \times 10^{-2}) = E_a/2.303 \times 8.314 [1/300 - 1/310]</math></p> <p><math>\log 2 = E_a/2.303 \times 8.314 [1/300 - 1/310]</math></p> <p><math>E_a = 53598.59 \text{ J/mol}</math> or <math>53.6 \text{ kJ/mol}</math></p> <p style="text-align: center;"><b>OR</b></p> <p>(a)(i) Rate becomes 4 times (ii) <math>2^{\text{nd}}</math> order</p> <p>b) <math>t_{1/2} = \frac{0.693}{k}</math></p> <p><math>23.1 \text{ min} = \frac{0.693}{k}</math></p> <p><math>k = 0.03 \text{ min}^{-1}</math></p> <p><math>k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}</math></p> <p><math>t = \frac{2.303}{0.03} \log \frac{100}{25}</math></p> <p><math>t = \frac{2.303}{0.03} \times 0.6021 \text{ min}</math></p> <p><math>t = 46.22 \text{ min}</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p>
25	<p>(i) X-X' bond in inter halogens is weaker than X-X in halogens</p> <p>(ii) High bond dissociation energy/ due to presence of triple bond.</p> <p>(iii) Because bond dissociation enthalpy decreases from <math>\text{NH}_3</math> to <math>\text{BiH}_3</math>.</p>	<p>1</p> <p>1</p> <p>1</p>

25	<p>b. (i)</p>  <p>(ii)</p>  <p><b>OR</b></p> <p>a) <math>\text{PH}_3</math>  <math>\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow 3\text{NaH}_2\text{PO}_2 + \text{PH}_3</math></p> <p>b) Helium</p> <p>c) Because bond dissociation energy of F-F bond is lower than that of Cl-Cl.</p> <p>d) <math>4\text{H}_3\text{PO}_3 \xrightarrow{\text{HEAT}} 3\text{H}_3\text{PO}_4 + \text{PH}_3</math></p> <p>e) <math>\text{PbS} + 4\text{O}_3 \rightarrow \text{PbSO}_4 + 4\text{O}_2</math></p>	1,1    1/2, 1/2 1 1 1 1 1
26	<p>(i)  <math>\text{CH}_3\text{CONH}_2 + \text{Br}_2 + 4\text{KOH} \rightarrow \text{CH}_3\text{NH}_2 + \text{K}_2\text{CO}_3 + 2\text{KBr} + 2\text{H}_2\text{O}</math></p> <p>(ii)  <math>\text{C}_6\text{H}_5\text{NH}_2 + \text{NaNO}_2 + 2\text{HCl} \xrightarrow{273-278\text{K}} \text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^- + \text{NaCl} + 2\text{H}_2\text{O}</math></p> <p>(iii)</p>  <p>b.(i) Add <math>\text{CHCl}_3</math> and alc KOH, aniline gives foul smell of isocyanide whereas N-methylaniline does not.</p> <p>(ii) When <math>(\text{CH}_3)_2\text{NH}</math> reacts with Benzene Sulphonyl Chloride (Hinsberg Reagent) gives ppt which is insoluble in alkali whereas <math>(\text{CH}_3)_3\text{N}</math> does not react with Hinsberg's Reagent.</p> <p>(Or any other correct distinguishing test)</p> <p><b>OR</b></p>	1  1  1  1  1

26	<p>a.</p> <p>(i)  (ii)  (iii) </p> <p>b. (i) <math>(\text{CH}_3)_3\text{N} &lt; \text{C}_2\text{H}_5\text{NH}_2 &lt; \text{C}_2\text{H}_5\text{OH}</math> (ii) p-nitroaniline &lt; aniline &lt; p-methylaniline</p>	1,1,1  1 1
----	--	---------------------