



Chemistry-Marking Scheme 2015

Chennai- 56/2/MT

| Q.No | Value points | Marks |
|------|---|-----------------|
| 1 | CH ₃ -CH ₂ -Br. | 1 |
| 2 | 1-methoxypropan-2-ol. | 1 |
| 3 | Dispersed phase – Solid , Dispersion medium – Liquid. | 1 |
| 4 | Due to incompletely filled d-orbitals in +2 oxidation state (ie., in Cu ²⁺ state.) | 1 |
| 5 | 3 Faraday / 3F | 1 |
| 6 | As per Raoult's law $p_A = x_A p_A^{\circ}$ $p_A = p_A^{\circ}(1 - x_B) = p_A^{\circ} - p_A^{\circ}x_B$ $(p_A^{\circ} - p_A) / p_A^{\circ} = x_B$ $\Delta p / p_A^{\circ} = x_B = \frac{W_B M_A}{M_B W_A}$ $M_B = \frac{W_B M_A}{(\Delta p / p_A^{\circ}) W_A}$ | 2 |
| 7 | (i) C ₆ H ₅ -NH ₂ < C ₆ H ₅ -NH-CH ₃ < CH ₃ -CH ₂ -NH ₂ . (ii) CH ₃ -NH-CH ₃ < CH ₃ -CH ₂ -NH ₂ < C ₂ H ₅ -OH. | 1 1 |
| 8 | Pentaamminecarbonatocobalt(III) chloride. Ionization isomerism | 1 1 |
| 8 | OR (i) [CuCl ₄] ²⁻ (ii) K ₂ [Zn(OH) ₄] | 1,1 |
| 9 | Rate constant is the proportionality constant that relates rate of reaction with concentration of reactants / Rate of the reaction when molar concentration of the reactant becomes unity. (i) Unit : time ⁻¹ or s ⁻¹ . (ii) Unit : L mol ⁻¹ time ⁻¹ or M ⁻¹ s ⁻¹ . | 1 ½ ½ |
| 10 |  i) (ii) | 1,1 |
| 11 | $\Delta T_b = K_b m$ $\Delta T_b = K_b (W_B \times 1000 / M_B \times W_A)$ $353.93 - 353.23 = 2.52 \times 1.5 \times 1000 / M_B \times 90$ $M_B = (2.52 \times 1.5 \times 1000) / (0.7 \times 90)$ = 60.0 g mol ⁻¹ . | 1 1 1 |

| 12 | (i) Because of $p\pi-p\pi$ multiple bonding in nitrogen (diatomic) which is absent in phosphorus (polymeric / polyatomic). | 1 | | | | | | | | | | | | |
|----|---|-----------------------------|-----------------|-----|-----|---|-------------------------|-----------------|---|--------------------------|---------------------------|--|---------------------------------------|--|
| | (ii) Because of decrease in tendency of sp^3 hybridisation from H_2O to H_2Te . | 1 | | | | | | | | | | | | |
| | (iii) Due to their smallest atomic sizes in respective periods, or due to the fact that they have only one electron less than the next noble gas configuration. | 1 | | | | | | | | | | | | |
| 13 | (i) $CH_3-CH(OH)-CH_3$ | 1 | | | | | | | | | | | | |
| | (ii) $CH_3-CH=CH-CH_3$ | 1 | | | | | | | | | | | | |
| | (iii) $p-Br-C_6H_4-CO-CH_3$ | 1 | | | | | | | | | | | | |
| 14 | (i) But-1,3-diene, Acrylonitrile; $CH_2=CH-CH=CH_2$, $CH_2=CH-CN$ | $\frac{1}{2} + \frac{1}{2}$ | | | | | | | | | | | | |
| | (ii) Phenol, Formaldehyde; C_6H_5OH , $HCHO$ | $\frac{1}{2} + \frac{1}{2}$ | | | | | | | | | | | | |
| | (iii) Tetrafluoroethylene; $CF_2=CF_2$ (Note: half mark for name/s and half mark for structure/s) | $\frac{1}{2} + \frac{1}{2}$ | | | | | | | | | | | | |
| 15 | (i) Gluconic acid or $COOH-(CHOH)_4-CH_2OH$ | 1 | | | | | | | | | | | | |
| | (ii) Peptide linkage or $-NH-CO-$ links | 1 | | | | | | | | | | | | |
| | (iii) | 1 | | | | | | | | | | | | |
| | <table border="1"> <thead> <tr> <th>s.no</th> <th>DNA</th> <th>RNA</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Sugar is 2-deoxy ribose</td> <td>Sugar is ribose</td> </tr> <tr> <td>2</td> <td>Double helical structure</td> <td>Single stranded structure</td> </tr> <tr> <td></td> <td colspan="2">(or any other one correct difference)</td> </tr> </tbody> </table> | | s.no | DNA | RNA | 1 | Sugar is 2-deoxy ribose | Sugar is ribose | 2 | Double helical structure | Single stranded structure | | (or any other one correct difference) | |
| | s.no | | DNA | RNA | | | | | | | | | | |
| 1 | Sugar is 2-deoxy ribose | | Sugar is ribose | | | | | | | | | | | |
| 2 | Double helical structure | Single stranded structure | | | | | | | | | | | | |
| | (or any other one correct difference) | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | |
| 16 | (a)(i) d^2sp^3 ; Octahedral | $\frac{1}{2} + \frac{1}{2}$ | | | | | | | | | | | | |
| | (ii) sp^3 ; Tetrahedral | $\frac{1}{2} + \frac{1}{2}$ | | | | | | | | | | | | |
| | (b) 'en', forms chelate. | $\frac{1}{2} + \frac{1}{2}$ | | | | | | | | | | | | |
| 17 | (i) Anion vacancies occupied by free electrons in alkali metal halides, (when they have metal excess defects) are called F-centre. | 1 | | | | | | | | | | | | |
| | (ii) When Si or Ge is doped with a trivalent impurity then electron vacancies are created called positive holes which impart electrical conduction. They are called p-type semiconductors. | 1 | | | | | | | | | | | | |
| | (iii) Ferrimagnetism is observed when the magnetic moments are aligned in parallel and antiparallel way in unequal numbers in a substance leading to small net permanent magnetic moment. | 1 | | | | | | | | | | | | |
| 18 | $\log (k_2 / k_1) = (E_a / 2.303R) (T_2 - T_1) / T_1 T_2$ | 1 | | | | | | | | | | | | |
| | $\log [(8 \times 10^{-2}) / (2 \times 10^{-2})] = 20 E_a / 2.303 \times 8.314 \times 300 \times 320$ | 1 | | | | | | | | | | | | |
| | $E_a = [\log(4) \times 2.303 \times 8.314 \times 300 \times 320] / 20$ | 1 | | | | | | | | | | | | |
| | $E_a = 55336.8 \text{ J mol}^{-1} = 55.34 \text{ kJ mol}^{-1}$. | 1 | | | | | | | | | | | | |
| 19 | (i) Due to intramolecular H-bonding in o-nitrophenol / | 1 | | | | | | | | | | | | |

| | | |
|----|---|--|
| | <p>p-nitrophenoxide is more stabilized than o-nitrophenoxide due to more delocalization of the negative charge.</p> <p>(ii) The mutual repulsion between bulky alkyl groups is stronger than the I.p-I.p electronic repulsions.</p> <p>(iii) CH₃ONa is not only nucleophile but also stronger base, thereby leads to elimination reaction of the alkyl halide.</p> | 1 1 1 |
| 20 | <p>(i) $C_6H_5NH_2 \xrightarrow{NaNO_2 + HCl / 278K} C_6H_5N_2Cl \xrightarrow{H_3PO_2 + H_2O} C_6H_6$</p> <p>(ii) $CH_3-CONH_2 \xrightarrow{KOH + Br_2} CH_3NH_2$</p> <p>(iii) $C_6H_5NO_2 \xrightarrow{Sn + HCl \text{ or } Fe + HCl} C_6H_5NH_2$</p> <p style="text-align: center;">OR</p> <p>(i) $C_2H_5NH_2 + CH_3COCl \xrightarrow{\text{pyridine}} C_2H_5-NHCOCH_3 + HCl$</p> <p>(ii) $C_2H_5NH_2 + C_6H_5SO_2Cl \longrightarrow C_2H_5NH-O_2SC_6H_5 + HCl$</p> <p>(iii) $C_2H_5NH_2 + CHCl_3 + KOH \longrightarrow C_2H_5NC + KCl + H_2O$</p> | 1 1 1 1 1 1 1 |
| 21 | <p>(i) In a catalysis process when the reactants and catalyst occur in same phase, the process is called homogeneous catalysis.</p> <p>(ii) The process of settling of colloidal particles forming precipitate is called coagulation.</p> <p>(iii) Polymeric substances or macromolecules when added to suitable solvents form solutions in which the size of the macromolecules may be in colloidal range. Such colloids are known as macromolecular colloids.</p> | 1 1 1 |
| 22 | <p>(i) The principle of zone refining is that the impurities are more soluble in the melt of metal than in solid state of the metal.</p> <p>(ii) As leaching agent, thereby oxidizing the metal into soluble cyanocomplex / [Au(CN)₂]⁻.</p> <p>(iii) Wrought iron</p> | 1 1 1 |
| 23 | <p>(i) Social awareness, Health conscious, Caring, empathy, concern. (or any other two values)</p> <p>(ii) (ii) Cartoon display / street display/poster making (or any other correct answer)</p> <p>(iii) Wrong choice and over dose may be harmful.</p> <p>(iv) Saccharin, Aspartame (or any other example)</p> | 1 1 1 $\frac{1}{2} + \frac{1}{2}$ |
| 24 | <p>(a)</p> <p>(i) Ce⁴⁺ gets reverted to 3+ oxidation state in aqueous medium hence is a good oxidizing agent / Ce is more stable in +3 oxidation state.</p> | 1 |

| | | |
|----|--|---|
| 24 | <p>(ii) Due to very strong metal-metal bonding (involving large no. of electrons of the d-orbitals)</p> <p>(iii) Mn has maximum no. of unpaired electrons in 3d-orbitals.</p> <p>(b)(i) $2\text{MnO}_4^- + 6\text{H}^+ + 5\text{NO}_2^- \longrightarrow 2\text{Mn}^{2+} + 5\text{NO}_3^- + 3\text{H}_2\text{O}$</p> <p>(ii) $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{Fe}^{2+} \longrightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$</p> <p style="text-align: center;">OR</p> <p>(a) (i) Due to d-d transitions (involving absorption of energy in visible range) / unpaired electrons in d- orbitals.</p> <p>(ii) Because Cr is more stable in +3 oxidation state.</p> <p>(iii) Due to stability of $5f^0$, $5f^7$, $5f^{14}$ / very small energy difference / comparable energy among 5f, 6d, and 7s orbitals.</p> <p>(b) The overall decrease in atomic and ionic radii from La to Lu (due to poor shielding effect of 4f electrons) is called Lanthanoid contraction. Common oxidation state of Lanthanoids is +3.</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1+1</p> |
| 25 | <p>(a) A is $\text{C}_6\text{H}_5\text{CHO}$; B & C / C & B are $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$ & $\text{C}_6\text{H}_5\text{COONa}$</p> <p>D is $\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{CH}_3$</p> <p>(b) (i) $\text{C}_6\text{H}_5\text{-CO-CH}_3$ forms yellow coloured CHI_3 on heating with I_2+KOH but $\text{C}_6\text{H}_5\text{-CO-CH}_2\text{-CH}_3$ does not / equation form.</p> <p>(ii) With neutral FeCl_3, phenol gives violet coloration but benzoic acid does not. (any other suitable test).</p> <p>(c) </p> <p style="text-align: center;">OR</p> <p>(a) (i) $\text{CH}_3\text{CH}(\text{OH})\text{CN}$</p> <p>(ii) $\text{CH}_3\text{CH}=\text{N-NH}_2$</p> <p>(iii) $\text{CH}_3\text{CH}_2\text{OH}$</p> <p>(b) $\text{C}_6\text{H}_5\text{-CO-CH}_3 < \text{CH}_3\text{-CO-CH}_3 < \text{CH}_3\text{-CHO}$</p> <p>(c) CH_3CHO gives yellow precipitate of CHI_3 with $\text{I}_2 + \text{KOH}$ but $\text{CH}_3\text{CH}_2\text{CHO}$ does not/ equation form</p> | <p>$\frac{1}{2} \times 4$</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
| 26 | <p>$E_{\text{cell}} = (E_{\text{Ag}}^{\circ} - E_{\text{Ni}}^{\circ}) - (0.0591/n) \log[\text{Ni}^{2+}/(\text{Ag}^+)^2]$</p> <p>$= (0.80 + 0.25) - 0.02955 \log(10^{-2}/10^{-6})$</p> <p>$= 1.05 - 0.0178 = 1.0322 \text{ V}$</p> <p>$\Delta G = - n F E_{\text{cell}}$</p> <p>$= - 2 \times 96500 \times 1.0322$</p> | <p>1</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |

| | | |
|----|--|--|
| | $= -199214 \text{ J mol}^{-1} = -199.2 \text{ kJ mol}^{-1}$ | 1 |
| | OR | |
| 26 | <p>(a) Molar Conductivity (Λ_m) = 1000 K / C</p> $= (1000 \times 1.06 \times 10^{-2}) / 0.1$ $= 106 \text{ S cm}^{-2} \text{ mol}^{-1}.$ | $\frac{1}{2}$ $\frac{1}{2}$ |
| | <p>Deg. of dissociation (α) = Λ_m / Λ_m^0</p> $= 106 / (50.1 + 76.5)$ $= 0.8373$ | 1 $\frac{1}{2}$ |
| | <p>(b) Primary battery- non rechargeable whereas secondary battery is chargeable. Eg: primary battery-dry cell, mercury cell(any one) , secondary battery- lead storage battery, Ni-Cd battery(any one) (or any other correct example)</p> | $\frac{1}{2}, \frac{1}{2}$ $\frac{1}{2}, \frac{1}{2}$ |