















	(ii) $\text{CH}_3\text{CH}_2\text{Cl} + \text{AgNO}_2 \rightarrow \text{CH}_3\text{CH}_2\text{NO}_2 + \text{AgCl}$	1
	(iii) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{Br})\text{CH}_3 + \text{KOH (alc.)} \rightarrow \text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_3$	1
Q.12	(i) Stoichiometric defect	1
	(ii) Schottky defect e.g. NaCl (or any other example)	$\frac{1}{2} + \frac{1}{2}$
	(iii) Density of crystal decreases	1
Q.13	$\Lambda_m = \frac{1000 \times k}{M} \text{Scm}^2\text{mol}^{-1}$	$\frac{1}{2}$
	$\Lambda_m = \frac{1000 \times 5.25 \times 10^{-5}}{2.5 \times 10^{-4}} \text{Scm}^2\text{mol}^{-1}$	1
	$= 210 \text{Scm}^2\text{mol}^{-1}$	
	$\Lambda_m^0 \text{HCOOH} = \lambda^0 \text{HCOO}^- + \lambda^0 \text{H}^+$	
	$(50.5 + 349.5) \text{S cm}^2\text{mol}^{-1} = 400 \text{S cm}^2\text{mol}^{-1}$	$\frac{1}{2}$
	$\alpha = \Lambda_m / \Lambda_m^0$	
	$\alpha = 210 / 400 = 0.525$	1
Q.14	i) $(\text{CH}_3)_2\text{C}=\text{N}-\text{NH}_2$ ii)  / benzoic acid iii)  / m-bromobenzoic acid	1+1+1
Q.15	(a)	
	(i) Because Cu^+ undergoes disproportionation as $2\text{Cu}^+ \rightarrow \text{Cu} + \text{Cu}^{2+}$	1
	(ii) Because of small size of metal, high ionic charge and availability of vacant d-orbital.	1
	(b) $\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 3\text{NO}_2^- \rightarrow 2\text{Cr}^{3+} + 3\text{NO}_3^- + 4\text{H}_2\text{O}$ (Balanced equation only)	1
Q.16	(i) ethylene glycol $\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}$	$\frac{1}{2} + \frac{1}{2}$
	Terephthalic acid 	

	<p>(ii) 1,3- butadiene $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$</p> <p style="text-align: center;">$\text{CH} = \text{CH}_2$</p>  <p>Styrene</p> <p>(iii) Chloroprene $\text{CH}_2=\text{C}(\text{Cl})-\text{CH}=\text{CH}_2$</p> <p>(Note: Half mark for name/s and half mark for structure/s in each case)</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p>
Q.17	$p^0 - p = \frac{w_s \times M_{\text{solvent}}}{M_s \times W_{\text{solvent}}}, \quad s = \text{solute}$ $(32 - 31.84)/32 = 10 \times 18 / M_s \times 200$ $M_s = 180 \text{ g/mol}$	<p>1</p> <p>1</p> <p>1</p>
Q.18	<p>(i) Zone refining</p> <p>(ii) SiO_2 act as flux to remove the impurity of Iron oxide</p> <p>(iii) Depressants prevent one type of sulphide ore forming the froth with air bubbles.</p>	<p>1</p> <p>1</p> <p>1</p>
Q.19	<p>Physisorption : adsorbate is held by weak van der Waals' force non-specific It forms multimolecular layer</p> <p>Chemisorption : adsorbate molecules are held by strong forces like a chemical bond It is specific It forms unimolecular layer (or any correct three points)</p>	<p>1,1,1</p>
Q.20	<p>(i) Phenoxide ion is stabilized by resonance as compared to CH_3O^- / In phenol, oxygen acquires +ve charge due to resonance and releases H^+ ion easily whereas there is no resonance in methanol.</p> <p>(ii) Due to lone pair- lone pair repulsion on oxygen.</p> <p>(iii) $(\text{CH}_3)_3\text{C}^+$ is 3^0 carbo-cation which is more stable than CH_3^+ for $\text{S}_{\text{N}}1$ reaction.</p>	<p>1</p> <p>1</p> <p>1</p>
Q.21	<p>(i) Starch.</p> <p>(ii) α- Helix polypeptide chains are stabilized by intramolecular H-bonding whereas β- pleated sheet is stabilized by intermolecular H-bonding. (or any other difference)</p> <p>(iii) Pernicious anaemia</p>	<p>1</p> <p>1</p> <p>1</p>
Q.22	<p>(i) Hydration isomerism</p> <p>(ii) Electronic configuration $1s^2 2g^4$ / by diagram</p> <p>(iii) Hybridization is $sp^3 d^2$ and shape is octahedral.</p>	<p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>

Q.23	<p>(i) Social awareness ,Health conscious, Caring , empathy, concern .(or any other two values)</p> <p>(ii) Cartoon display / street play/poster making (or any other correct answer)</p> <p>(iii) Wrong choice and over dose may be harmful.</p> <p>(iv) Saccharin , Aspartame (or any other example)</p>	<p>½ , ½</p> <p>1</p> <p>1</p> <p>½ + ½</p>
Q.24	<p>A = </p> <p>B = </p> <p>C = </p> <p>D = </p> <p>E = </p> <p>OR</p> <p>Q24</p> <p>i) </p> <p>ii) </p> <p>iii) </p> <p>(b) $C_2H_5NH_2 < (C_2H_5)_3N < (C_2H_5)_2NH$</p> <p>(c) Add $CHCl_3$ and alc KOH , $C_6H_5-NH_2$ gives foul smell of isocyanide whereas $C_6H_5-NH-CH_3$ does not (or any other correct test)</p>	<p>1 x 5</p> <p>1,1,1</p> <p>1</p> <p>1</p>
Q.25	<p>(a) $[A]_0 = 0.10 \text{ mol/L}$ $[A] = 0.05 \text{ mol/L}$ at time $t = 10\text{s}$</p> $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $k = \frac{2.303}{10 \text{ s}} \log \frac{0.10}{0.05}$	<p>½</p>

	$k = 0.0693 \text{ s}^{-1}$	1
	$t = 20\text{s}$ $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$	
	$k = \frac{2.303}{20 \text{ s}} \log \frac{0.10}{0.025}$ $k = 0.0693 \text{ s}^{-1}$	1
	As the rate constant is same , it follows pseudo first order reaction.	$\frac{1}{2}$
	(b) Average rate of reaction = $-\Delta[R] / \Delta t$	$\frac{1}{2}$
	$= - [0.025 - 0.05 / 20 - 10]$	$\frac{1}{2}$
	$= 0.0025 \text{ mol L}^{-1}\text{s}^{-1}$	1
Q25	OR	
	(a)	1
	(i) Rate of reaction becomes 4 times	1
	(ii) Over all order of reaction = 2	
	(b) $t_{1/2} = \frac{0.693}{k}$	
	$30\text{min} = \frac{0.693}{k}$	
	$k = 0.0231\text{min}^{-1}$	1
	$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$	$\frac{1}{2}$
	$t = \frac{2.303}{0.0231} \log \frac{100}{10}$	$\frac{1}{2}$

	$t = \frac{2.303}{0.0231} \min$ $t = 99.7 \min$	1
Q.26	<p>(a) (i) Due to decrease in bond dissociation enthalpy from HF to HI , there is an increase in acidic character observed.</p> <p>(ii)Oxygen exists as diatomic O₂ molecule while sulphur as polyatomic S₈</p> <p>(iii)Due to non- availability of d orbitals</p> <p>(b)</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>i)</p> </div> <div style="text-align: center;">  <p>ii)</p> </div> </div>	1 1 1 1+1
Q26	OR	
	<p>(i) White Phosphorus, because it is less stable due to angular strain</p> <p>(ii)Nitrogen oxides emitted by supersonic jet planes are responsible for depletion of ozone layer. Or $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$</p> <p>(iii)due to small size of F, large inter electronic repulsion / electron- electron repulsion among the lone pairs of fluorine</p> <p>(iv) Helium</p> <p>(v) $\text{XeF}_2 + \text{PF}_5 \rightarrow [\text{XeF}]^+ [\text{PF}_6]^-$</p>	1/2, 1/2 1 1 1 1