






Class: XII
Chemistry
Marking Scheme 2018-19

Time allowed: 3 Hours



Maximum Marks: 70


Q No	SECTION A	Marks
1.	On heating ZnO , it loses oxygen and there is excess of Zn^{2+} ions in the crystal.	1
	OR	
	When silicon is doped with phosphorous (group 15 element), the increase in conductivity is due to the delocalised negatively charged electrons.	1
2.	Associated colloids	1
3.	$t_{2g}^3 e_g^3$	1
	OR	
	The orbital splitting energies, Δ_t are not sufficiently large for forcing pairing of electrons in the tetrahedral coordination entity formation.	1
4.	B	1
5.	<i>Poly β – hydroxybutyrate – co – β – hydroxy valerate</i>	1
SECTION B		
6.	$HBr \rightarrow H^+ + Br^-$ $i = 1 - \alpha + n\alpha$ $n = 2$ $i = 1 + \alpha$ $\Delta T_f = iK_f m$ $\Delta T_f = (1 + \alpha) 1.86 \times \frac{8.1}{81} \times \frac{1000}{100}$ $\Delta T_f = 3.53$ $T_f^0 = 0^0 C$ $\Delta T_f = T_f^0 - T_f'$ $T_f' = -3.534^0 C$	 1/2 1/2 1/2 1/2 1/2 1/2
	OR	
	<i>Mole fraction of water, $\chi_{H_2O} = 0.88$</i> <i>Mole fraction of ethanol, $\chi_{C_2H_5OH} = 1 - 0.88$</i> $= 0.12$ $\chi_{C_2H_5OH} = \frac{n_2}{n_1 + n_2} \dots\dots\dots(1)$	1/2

	<p>n_2 = number of moles of ethanol. n_1 = number of moles of water. Molality of ethanol means the number of moles of ethanol present in 1000 g of water.</p> $n_1 = \frac{1000}{18} = 55.5 \text{ moles}$ <p>Substituting the value of n_1 in equation (1)</p> $\frac{n_2}{55.5 + n_2} = 0.12$ <p>$n_2 = 7.57 \text{ moles}$ Molality of ethanol ($\text{C}_2\text{H}_5\text{OH}$) = 7.57 m</p> <p>Alternatively,</p> <p>Mole fraction of water = 0.88 Mole fraction of ethanol = 1 - 0.88 = 0.12 Therefore 0.12 moles of ethanol are present in 0.88 moles of water. Mass of water = 0.88 x 18 = 15.84 g of water. Molality = number of moles of solute (ethanol) present in 1000 g of solvent (water)</p> $= 12 \times 1000 / 15.84$ $= 7.57 \text{ m}$ Molality of ethanol ($\text{C}_2\text{H}_5\text{OH}$) = 7.57 m	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
7.(a)	Reaction : Hell-Volhard-Zelinsky reaction. IUPAC : 2-Bromopropanoic acid.	$\frac{1}{2}$
(b)	Reaction : Rosenmund reduction reaction. IUPAC : Benzaldehyde.	$\frac{1}{2}$
	OR	
	i) 2-Methylbut-2-enal	$\frac{1}{2}$
	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3\text{-CH=CH-CHO} \end{array}$	$\frac{1}{2}$
	ii) Pent-2-enal	$\frac{1}{2}$
	$\text{CH}_3\text{-CH}_2\text{-CH=CH-CHO}$	$\frac{1}{2}$
8.(a)	Tert-butyl peroxide acts as a free radical generating initiator(catalyst)	1
(b)	$\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$	$\frac{1}{2}$
	$\text{C}_6\text{H}_5 - \text{CH} = \text{CH}_2$	$\frac{1}{2}$



9.	<p>Step 1 : Formation of protonated alcohol.</p>  <p>Ethanol</p> <p>Protonated ethanol (Ethyl oxonium ion)</p> <p>Step 2 : Formation of carbocation.</p>  <p>Step 3: Formation of ethane by elimination of a proton.</p>  <p>Ethene</p>	<p>1/2</p> <p>1</p> <p>1/2</p>
10.	<p>(i) Zero order reaction (ii) Slope represents $-k$; Intercept represents $[R]_0$ (iii) $\text{mol L}^{-1} \text{s}^{-1}$</p>	<p>1/2 1/2+1/2 1/2</p>
11.(a)	 <p>XeO₃</p>	1
(b)	 <p>H₃PO₃</p>	1
12.(i)	<p>Since $\text{Ti}^{4+}/\text{Ti}^{3+}$ has lower reduction potential than $\text{Fe}^{3+}/\text{Fe}^{2+}$, it cannot be reduced in comparison with $\text{Fe}^{3+}/\text{Fe}^{2+}$ ions. Hence Ti^{4+} cannot oxidise Fe^{2+} to Fe^{3+}.</p>	1 1/2
(ii)	<p>As the value of reduction potential increases the stability of +2 oxidation increases. Therefore correct order of stability is $\text{Cr}^{3+}/\text{Cr}^{2+} < \text{Fe}^{3+}/\text{Fe}^{2+} < \text{Mn}^{3+}/\text{Mn}^{2+}$</p>	1 1/2

SECTION C		
13.	$r = \frac{\sqrt{3}}{4} a$ $143.1 = \frac{\sqrt{3}}{4} a$ $a = 330.4 \text{ pm}$ $\rho = \frac{zM}{a^3 N_A}$ $\rho = \frac{2 \times 93}{(330.4 \times 10^{-10})^3 \times 6.023 \times 10^{23}}$ $\rho = 8.58 \text{ g/cm}^3$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$
14.(a)	Molecules of benzoic acid dimerise in benzene, the number of particles are reduced.	1
(b)	The intermolecular interactions between ethanol and acetone are weaker/ the escaping tendency of ethanol and acetone molecules increases on mixing / the vapour pressure increases.	1
(c)	Due to osmosis, a bacterium on fruit loses water, shrivels and dies.	1
15.	<p style="text-align: center;"> $\begin{array}{c} \text{CH}_3 \\ \\ \text{A : CH}_3\text{-CH-CH}_2\text{-OH} \end{array}$ </p> <p style="text-align: center;"> $\begin{array}{c} \text{CH}_3 \\ \\ \text{B : CH}_3\text{-CH-COOH} \end{array}$ </p> <p style="text-align: center;"> $\begin{array}{c} \text{CH}_3 \\ \\ \text{C : CH}_3\text{-CH} = \text{CH}_2 \end{array}$ </p> <p style="text-align: center;"> $\begin{array}{c} \text{CH}_3 \\ \\ \text{D : CH}_3\text{-CH-CH}_3 \\ \\ \text{OH} \end{array}$ </p> <p>A and D are position isomers.</p>	$\mathbf{1}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

16.	<p>$C_6H_5CH_2Cl$ will undergo S_N1 reaction faster. The carbocation formed by $C_6H_5CH_2Cl$ gets stabilized through resonance. Greater the stability of carbocation, greater will be its ease of formation from the respective halide.</p> 	<p>$\frac{1}{2}$ $\frac{1}{2}$ 1 1</p>
OR		
	<p style="text-align: center;"> $\begin{array}{c} Cl \\ \\ CH_3-CH-CH_2Cl \end{array}$ </p> <p style="text-align: center;">$CH_2Cl-CH_2-CH_2Cl$</p> <p style="text-align: center;">$CH_3-CH_2-CHCl_2$</p> <p style="text-align: center;"> $\begin{array}{c} Cl \\ \\ CH_3-C-CH_3 \\ \\ Cl \end{array}$ </p> <p>The following isomer will exhibit enantiomerism:</p> <p style="text-align: center;"> $\begin{array}{c} Cl \\ \\ CH_3-CH-CH_2Cl \end{array}$ </p> <p>IUPAC name: 1,2-Dichloropropane.</p>	<p>$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$</p>
17.(a)		1
(b)	$HOOC-CH_2-CH_2-CH_2-CH_2-COOH$	1

(c)		1
18.(i)	It is unstable at cooking temperature.	1
(ii)	Excessive hydrogencarbonate can make the stomach alkaline and trigger the production of even more acid. Metal hydroxides being insoluble do not increase the pH above neutrality.	1
(iii)	Aspirin has anti blood clotting action.	1
19.(a)	Amylopectin.	1
(b)	C- 2	1
(c)	Two peptide linkages.	1
OR		
	(1) Glucose does not give 2,4- DNP test.	1
	(2) Glucose does not give Schiff's test.	1
	(3) The pentaacetate of glucose does not react with hydroxylamine.	1
	(4) Glucose does not form the hydrogensulphite addition product with NaHSO ₃ . (Any three points)	
20.	$2N_2O_5(g) \rightarrow 2N_2O_4(g) + O_2(g)$ <p>At t = 0 0.5 atm 0 atm 0 atm</p> <p>At time t 0.5 - 2x atm 2x atm x atm</p> $P_t = P_{N_2O_5} + P_{N_2O_4} + P_{O_2}$ $= (0.5 - 2x) + 2x + x = 0.5 + x$ $x = p_t - 0.5$ $P_{N_2O_5} = 0.5 - 2x$ $= 0.5 - 2(p_t - 0.5)$ $= 1.5 - 2p_t$ <p>At t = 100 s ; p_t = 0.512 atm</p> $P_{N_2O_5} = 1.5 - 2 \times 0.512 = 0.476 \text{ atm}$ $k = \frac{2.303}{t} \log \frac{p_i}{p_A}$ $k = \frac{2.303}{100s} \log \frac{0.5 \text{ atm}}{0.476 \text{ atm}}$ $k = \frac{2.303}{100s} \times 0.0216 = 4.98 \times 10^{-4} s^{-1}$	<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1</p>
OR		1
The Arrhenius equation: $k = Ae^{-E_a/RT}$		
Taking log on both sides: $\log k = \log A - \frac{E_a}{2.303RT}$		

	<p>For reaction (i) $\log k_1 = \log A - \frac{E_a(1)}{2.303RT}$</p> <p>For reaction (ii) $\log k_2 = \log A - \frac{E_a(2)}{2.303RT}$</p> <p>Subtracting (i) from (ii)</p> $\log \frac{k_1}{k_2} = \frac{E_a(1) - E_a(2)}{2.303RT}$ $= \log \frac{k_1}{k_2} = \frac{24.9 \times 1000}{2.303 \times 8.3 \times 300} = 4.342$ $\frac{k_1}{k_2} = \text{anti log}(4.342) = 2.198 \times 10^4$	1 1
21.		
(a)	Negative charge is developed on the sol. Sol is represented as AgI / I^-	$\frac{1}{2}$ $\frac{1}{2}$
(b)	Adsorption of reactants on the solid surface of the catalysts increases the rate of reaction.	1
(c)	Na_3PO_4 Hardy-Schulze rule	$\frac{1}{2}$ $\frac{1}{2}$
22.		
(a)	Leached gold complex is treated with Zinc and gold is recovered by displacement method $2Au[(CN)_2]^- (aq) + Zn(s) \rightarrow 2Au(s) + [Zn(CN)_4]^{2-} (aq)$	$\frac{1}{2}$ $\frac{1}{2}$
(b)	Zirconium iodide is decomposed on a tungsten filament; electrically heated to 1800 K. Pure Zr metal is deposited on the filament. $ZrI_4 \rightarrow Zr + I_2$	$\frac{1}{2}$ $\frac{1}{2}$
(c)	Silica is added to the ore and heated. It helps to slag off iron oxide as iron silicate $FeO + SiO_2 \rightarrow FeSiO_3(\text{slag})$	$\frac{1}{2}$ $\frac{1}{2}$
OR		
(a)	NaCN is used as depressants to separate two sulphide ores (ZnS and PbS) in Froth Floatation Method.	(1)
(b)	Carbon monoxide forms a volatile complex of nickel, nickel tetracarbonyl.	(1)
(c)	Coke is used as a reducing agent to reduce zinc oxide to zinc.	(1)
23.		
(a)	$Co^{2+} : [Ar]3d^7$ $Sc^{3+} : [Ar]3d^0$ $Cr^{3+} : [Ar]3d^3$ Co^{2+} and Cr^{3+} have unpaired electrons. Thus, they are coloured in aqueous solution. Sc^{3+} has no unpaired electron. Thus it is colourless.	1
(b)	Metal copper has high enthalpy of atomisation and enthalpy of ionisation. Therefore the high energy required to convert $Cu(s)$ to $Cu^{2+}(aq)$ is not balanced by its hydration enthalpy.	1

(c)	Due to lanthanoid contraction the size of lanthanoid ion decreases regularly with increase in atomic size. Thus covalent character between lanthanoid ion and OH^- increases from La^{3+} to Lu^{3+} . Thus the basic character of hydroxides decreases from $\text{La}(\text{OH})_3$ to $\text{Lu}(\text{OH})_3$	1						
24. (a) (b) (c)	Isomer A: $[\text{Cr}(\text{NH}_3)_4 \text{BrCl}] \text{Cl}$ Isomer B: $[\text{Cr}(\text{NH}_3)_4 \text{Cl}_2] \text{Br}$ Hybridisation of Cr in isomer A and B is d^2sp^3 . Number of unpaired electrons in $\text{Cr}^{3+}(3d^3)$ is 3 Magnetic moment = $\sqrt{n(n+2)}$ $= \sqrt{3(3+2)} = 3.87 \text{ BM}$ (deduct half mark for wrong unit/unit not written)	$\frac{1}{2}$ $\frac{1}{2}$ 1 1						
25.(a)	A = AgNO_2 B =  C =  D = $\text{CHCl}_3 + \text{KOH}$, Heat.	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$						
(b)	<table border="1"> <thead> <tr> <th>Experiment</th> <th>Aniline</th> <th>Benzylamine</th> </tr> </thead> <tbody> <tr> <td>Azo dye test: Dissolve the amine in HCl, cool it and then add cold aqueous solution of NaNO_2 and then solution of β-naphthol</td> <td>A brilliant orange red dye is observed.</td> <td>No dye is formed.</td> </tr> </tbody> </table>	Experiment	Aniline	Benzylamine	Azo dye test: Dissolve the amine in HCl, cool it and then add cold aqueous solution of NaNO_2 and then solution of β -naphthol	A brilliant orange red dye is observed.	No dye is formed.	1
Experiment	Aniline	Benzylamine						
Azo dye test: Dissolve the amine in HCl, cool it and then add cold aqueous solution of NaNO_2 and then solution of β -naphthol	A brilliant orange red dye is observed.	No dye is formed.						
(i)								
(ii)	<table border="1"> <thead> <tr> <th>Experiment</th> <th>Methylamine</th> <th>Dimethylamine</th> </tr> </thead> <tbody> <tr> <td>Carbylamine test: To the organic compound add chloroform and ethanolic potassium hydroxide and heat (or any other suitable test)</td> <td>A foul smelling substance (isocyanide)</td> <td>No reaction.</td> </tr> </tbody> </table>	Experiment	Methylamine	Dimethylamine	Carbylamine test: To the organic compound add chloroform and ethanolic potassium hydroxide and heat (or any other suitable test)	A foul smelling substance (isocyanide)	No reaction.	1
Experiment	Methylamine	Dimethylamine						
Carbylamine test: To the organic compound add chloroform and ethanolic potassium hydroxide and heat (or any other suitable test)	A foul smelling substance (isocyanide)	No reaction.						

(c)	A = CH ₃ CH ₂ CH ₂ NH ₂ B = CH ₃ CH ₂ CH ₂ OH	1/2 1/2
OR		
(a)(i)	In strongly acidic medium, aniline is protonated to form the anilium ion which is meta directing.	1
(ii)	Aryl halides do not undergo nucleophilic substitution with the anion formed by phthalimide.	1
(b)(i)	$\text{CH}_3\text{-COOH} \xrightarrow{\text{NH}_3, \Delta} \text{CH}_3\text{-CO-NH}_2 \xrightarrow{\text{NaOH} + \text{Br}_2} \text{CH}_3\text{NH}_2$	1
(ii)		1
(c)	p- Nitroaniline < Aniline < p- Toludine.	1
26.(a)	E _{cell} decreases.	1
(b)	Anode: Cl ₂ ↑ Cathode : H ₂ ↑	1/2 1/2
(c)	$\text{Cu}^{2+}(\text{aq.}) + \text{Ni}(\text{s}) \rightarrow \text{Ni}^{2+}(\text{aq.}) + \text{Cu}(\text{s})$ $E_{\text{cell}}^0 = E_{\text{cathode}}^0 - E_{\text{anode}}^0$ $E_{\text{cell}}^0 = 0.34 - (-0.25)$ $E_{\text{cell}}^0 = 0.59\text{V}$ $E_{\text{cell}} = E_{\text{cell}}^0 - \frac{2.303RT}{nF} \log \frac{[\text{Ni}^{2+}]}{[\text{Cu}^{2+}]}$ $E_{\text{cell}} = 0.59 - \frac{0.059}{2} \log \frac{[0.01]}{[0.1]}$ $E_{\text{cell}} = 0.6195\text{V}$	1/2 1/2 1/2 + 1/2
OR		
(a)	$\Lambda_{m(\text{CaCl}_2)}^0 = \lambda_{\text{Ca}^{2+}}^0 + 2\lambda_{\text{Cl}^-}^0$	1
(b)	Conductivity of NaCl decreases on dilution as the number of ions per unit volume decreases.	1

	Whereas molar conductivity of NaCl increases on dilution as on dilution the interionic interactions are overcome and ions are free to move.	1
(c)	$G^* = \kappa R$	1/2
	$\kappa = \frac{1.29}{100} = 0.0129 \text{ S cm}^{-1}$	1/2
	$\Lambda_m = \frac{1000 \kappa}{C}$	1/2
	$\Lambda_m = \frac{1000 \times 0.0129}{0.1}$	
	$\Lambda_m = 129 \text{ S cm}^2 \text{ mol}^{-1}$	1/2
27.		
(a)	(i) S atom in SF ₄ is not sterically protected as it is surrounded by only four F atoms, so attack of water molecules can take place easily. In contrast, S atom in SF ₆ is protected by six F atoms. Thus attack by water molecules cannot take place easily.	1
	(ii) Chlorine water produces nascent oxygen (causes oxidation) which is responsible for bleaching action. $Cl_2 + H_2O \rightarrow 2HCl + [O]$	1
	(iii) Due to inert pair effect Bi(V) can accept a pair of electrons to form more stable Bi(III). (+3 oxidation state of Bi is more stable than its +5 oxidation state).	1
(b)	(i) Phosphorus undergoes disproportionation reaction to form phosphine gas. $P_4 + 3NaOH + 3H_2O \rightarrow PH_3 + 3NaH_2PO_2$	1
	(ii) On partial hydrolysis, XeF ₆ gives oxyfluoride XeOF ₄ and HF. $XeF_6 + H_2O \rightarrow XeOF_4 + 2HF$	1
	OR	
(a)	N. Bartlett first prepared a red compound O ₂ ⁺ PtF ₆ ⁻ . He then realised that the first ionisation enthalpy of molecular oxygen was almost identical with Xenon. So he carried out reaction between Xe and PtF ₆ .	1
(b)	(i) I ₂ < F ₂ < Br ₂ < Cl ₂	1
	(ii) NH ₃ > PH ₃ > AsH ₃ > SbH ₃ > BiH ₃	1
(c)	(i) $2NaOH + Cl_2 \rightarrow NaCl + NaOCl + H_2O$	1
	(ii) $2Fe^{3+} + SO_2 + 2H_2O \rightarrow 2Fe^{2+} + SO_4^{2-} + 4H^+$	1
