

# COMPLEX NUMBERS AND QUADRATIC EQUATIONS

## 5.1 Overview

We know that the square of a real number is always non-negative e.g.  $(4)^2 = 16$  and  $(-4)^2 = 16$ . Therefore, square root of 16 is  $\pm 4$ . What about the square root of a negative number? It is clear that a negative number can not have a real square root. So we need to extend the system of real numbers to a system in which we can find out the square roots of negative numbers. Euler (1707 - 1783) was the first mathematician to introduce the symbol  $i$  (iota) for positive square root of  $-1$  i.e.,  $i = \sqrt{-1}$ .

### 5.1.1 Imaginary numbers

Square root of a negative number is called an imaginary number, for example,

$$\sqrt{-9} = \sqrt{-1} \sqrt{9} = i3, \quad \sqrt{-7} = \sqrt{-1} \sqrt{7} = i\sqrt{7}$$

### 5.1.2 Integral powers of $i$

$$i = \sqrt{-1}, \quad i^2 = -1, \quad i^3 = i^2 i = -i, \quad i^4 = (i^2)^2 = (-1)^2 = 1.$$

To compute  $i^n$  for  $n > 4$ , we divide  $n$  by 4 and write it in the form  $n = 4m + r$ , where  $m$  is quotient and  $r$  is remainder ( $0 \leq r < 4$ )

$$\text{Hence } i^n = i^{4m+r} = (i^4)^m \cdot (i)^r = (1)^m (i)^r = i^r$$

$$\text{For example, } (i)^{39} = i^{4 \times 9 + 3} = (i^4)^9 \cdot (i)^3 = i^3 = -i$$

$$\begin{aligned} \text{and } (i)^{-435} &= i^{-(4 \times 108 + 3)} = (i)^{-(4 \times 108)} \cdot (i)^{-3} \\ &= \frac{1}{(i^4)^{108}} \cdot \frac{1}{(i)^3} = \frac{i}{(i)^4} = i \end{aligned}$$

- (i) If  $a$  and  $b$  are positive real numbers, then

$$\sqrt{-a} \times \sqrt{-b} = \sqrt{-1} \sqrt{a} \times \sqrt{-1} \sqrt{b} = i\sqrt{a} \times i\sqrt{b} = -\sqrt{ab}$$

- (ii)  $\sqrt{a} \cdot \sqrt{b} = \sqrt{ab}$  if  $a$  and  $b$  are positive or at least one of them is negative or zero. However,  $\sqrt{a} \sqrt{b} \neq \sqrt{ab}$  if  $a$  and  $b$ , both are negative.

**5.1.3 Complex numbers**

- (a) A number which can be written in the form  $a + ib$ , where  $a, b$  are real numbers and  $i = \sqrt{-1}$  is called a complex number.
- (b) If  $z = a + ib$  is the complex number, then  $a$  and  $b$  are called real and imaginary parts, respectively, of the complex number and written as  $\text{Re}(z) = a, \text{Im}(z) = b$ .
- (c) Order relations “greater than” and “less than” are not defined for complex numbers.
- (d) If the imaginary part of a complex number is zero, then the complex number is known as purely real number and if real part is zero, then it is called purely imaginary number, for example, 2 is a purely real number because its imaginary part is zero and  $3i$  is a purely imaginary number because its real part is zero.

**5.1.4 Algebra of complex numbers**

- (a) Two complex numbers  $z_1 = a + ib$  and  $z_2 = c + id$  are said to be equal if  $a = c$  and  $b = d$ .
- (b) Let  $z_1 = a + ib$  and  $z_2 = c + id$  be two complex numbers then  $z_1 + z_2 = (a + c) + i(b + d)$ .

**5.1.5 Addition of complex numbers satisfies the following properties**

- As the sum of two complex numbers is again a complex number, the set of complex numbers is closed with respect to addition.
- Addition of complex numbers is commutative, i.e.,  $z_1 + z_2 = z_2 + z_1$
- Addition of complex numbers is associative, i.e.,  $(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3)$
- For any complex number  $z = x + iy$ , there exist 0, i.e.,  $(0 + 0i)$  complex number such that  $z + 0 = 0 + z = z$ , known as identity element for addition.
- For any complex number  $z = x + iy$ , there always exists a number  $-z = -a - ib$  such that  $z + (-z) = (-z) + z = 0$  and is known as the additive inverse of  $z$ .

**5.1.6 Multiplication of complex numbers**

Let  $z_1 = a + ib$  and  $z_2 = c + id$ , be two complex numbers. Then

$$z_1 \cdot z_2 = (a + ib)(c + id) = (ac - bd) + i(ad + bc)$$

- As the product of two complex numbers is a complex number, the set of complex numbers is closed with respect to multiplication.
- Multiplication of complex numbers is commutative, i.e.,  $z_1 \cdot z_2 = z_2 \cdot z_1$
- Multiplication of complex numbers is associative, i.e.,  $(z_1 \cdot z_2) \cdot z_3 = z_1 \cdot (z_2 \cdot z_3)$

4. For any complex number  $z = x + iy$ , there exists a complex number 1, i.e.,  $(1 + 0i)$  such that  
 $z \cdot 1 = 1 \cdot z = z$ , known as identity element for multiplication.
5. For any non zero complex number  $z = x + iy$ , there exists a complex number  $\frac{1}{z}$  such that  $z \cdot \frac{1}{z} = \frac{1}{z} \cdot z = 1$ , i.e., multiplicative inverse of  $a + ib = \frac{1}{a+ib} = \frac{a-ib}{a^2+b^2}$ .
6. For any three complex numbers  $z_1, z_2$  and  $z_3$ ,
- $$z_1 \cdot (z_2 + z_3) = z_1 \cdot z_2 + z_1 \cdot z_3$$
- and
- $$(z_1 + z_2) \cdot z_3 = z_1 \cdot z_3 + z_2 \cdot z_3$$
- i.e., for complex numbers multiplication is distributive over addition.

**5.1.7** Let  $z_1 = a + ib$  and  $z_2 (\neq 0) = c + id$ . Then

$$z_1 \div z_2 = \frac{z_1}{z_2} = \frac{a+ib}{c+id} = \frac{(ac+bd)}{c^2+d^2} + i \frac{(bc-ad)}{c^2+d^2}$$

### 5.1.8 Conjugate of a complex number

Let  $z = a + ib$  be a complex number. Then a complex number obtained by changing the sign of imaginary part of the complex number is called the conjugate of  $z$  and it is denoted by  $\bar{z}$ , i.e.,  $\bar{z} = a - ib$ .

Note that additive inverse of  $z$  is  $-a - ib$  but conjugate of  $z$  is  $a - ib$ .

We have :

- $\overline{(\bar{z})} = z$
- $z + \bar{z} = 2 \operatorname{Re}(z)$ ,  $z - \bar{z} = 2i \operatorname{Im}(z)$
- $z = \bar{z}$ , if  $z$  is purely real.
- $z + \bar{z} = 0 \Leftrightarrow z$  is purely imaginary
- $z \cdot \bar{z} = \{\operatorname{Re}(z)\}^2 + \{\operatorname{Im}(z)\}^2$ .
- $\overline{(z_1 + z_2)} = \bar{z}_1 + \bar{z}_2$ ,  $\overline{(z_1 - z_2)} = \bar{z}_1 - \bar{z}_2$
- $\overline{(z_1 \cdot z_2)} = (\bar{z}_1) (\bar{z}_2)$ ,  $\overline{\left(\frac{z_1}{z_2}\right)} = \frac{(\bar{z}_1)}{(\bar{z}_2)}$  ( $\bar{z}_2 \neq 0$ )

### 5.1.9 Modulus of a complex number

Let  $z = a + ib$  be a complex number. Then the positive square root of the sum of square of real part and square of imaginary part is called modulus (absolute value) of  $z$  and it is denoted by  $|z|$  i.e.,  $|z| = \sqrt{a^2 + b^2}$

In the set of complex numbers  $z_1 > z_2$  or  $z_1 < z_2$  are meaningless but

$$|z_1| > |z_2| \text{ or } |z_1| < |z_2|$$

are meaningful because  $|z_1|$  and  $|z_2|$  are real numbers.

### 5.1.10 Properties of modulus of a complex number

- $|z| = 0 \Leftrightarrow z = 0$  i.e.,  $\operatorname{Re}(z) = 0$  and  $\operatorname{Im}(z) = 0$
- $|z| = |\bar{z}| = |-z|$
- $-|z| \leq \operatorname{Re}(z) \leq |z|$  and  $-|z| \leq \operatorname{Im}(z) \leq |z|$
- $z \bar{z} = |z|^2$ ,  $|z^2| = |\bar{z}|^2$
- $|z_1 z_2| = |z_1| \cdot |z_2|$ ,  $\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$  ( $z_2 \neq 0$ )
- $|z_1 + z_2|^2 = |z_1|^2 + |z_2|^2 + 2\operatorname{Re}(z_1 \bar{z}_2)$
- $|z_1 - z_2|^2 = |z_1|^2 + |z_2|^2 - 2\operatorname{Re}(z_1 \bar{z}_2)$
- $|z_1 + z_2| \leq |z_1| + |z_2|$
- $|z_1 - z_2| \geq |z_1| - |z_2|$
- $|az_1 - bz_2|^2 + |bz_1 + az_2|^2 = (a^2 + b^2)(|z_1|^2 + |z_2|^2)$

In particular:

$$|z_1 - z_2|^2 + |z_1 + z_2|^2 = 2(|z_1|^2 + |z_2|^2)$$

- As stated earlier multiplicative inverse (reciprocal) of a complex number  $z = a + ib$  ( $\neq 0$ ) is

$$\frac{1}{z} = \frac{a - ib}{a^2 + b^2} = \frac{\bar{z}}{|z|^2}$$

## 5.2 Argand Plane

A complex number  $z = a + ib$  can be represented by a unique point P ( $a, b$ ) in the cartesian plane referred to a pair of rectangular axes. The complex number  $0 + 0i$  represent the origin O ( $0, 0$ ). A purely real number  $a$ , i.e., ( $a + 0i$ ) is represented by the point ( $a, 0$ ) on  $x$ -axis. Therefore,  $x$ -axis is called real axis. A purely imaginary number

$ib$ , i.e.,  $(0 + ib)$  is represented by the point  $(0, b)$  on  $y$ -axis. Therefore,  $y$ -axis is called imaginary axis.

Similarly, the representation of complex numbers as points in the plane is known as **Argand diagram**. The plane representing complex numbers as points is called complex plane or Argand plane or Gaussian plane.

If two complex numbers  $z_1$  and  $z_2$  be represented by the points P and Q in the complex plane, then

$$|z_1 - z_2| = PQ$$

### 5.2.1 Polar form of a complex number

Let P be a point representing a non-zero complex number  $z = a + ib$  in the Argand plane. If OP makes an angle  $\theta$  with the positive direction of  $x$ -axis, then  $z = r(\cos\theta + i\sin\theta)$  is called the polar form of the complex number, where

$r = |z| = \sqrt{a^2 + b^2}$  and  $\tan\theta = \frac{b}{a}$ . Here  $\theta$  is called argument or amplitude of  $z$  and we write it as  $\arg(z) = \theta$ .

The unique value of  $\theta$  such that  $-\pi \leq \theta \leq \pi$  is called the principal argument.

$$\arg(z_1 \cdot z_2) = \arg(z_1) + \arg(z_2)$$

$$\arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$$

### 5.2.2 Solution of a quadratic equation

The equations  $ax^2 + bx + c = 0$ , where  $a, b$  and  $c$  are numbers (real or complex,  $a \neq 0$ ) is called the general quadratic equation in variable  $x$ . The values of the variable satisfying the given equation are called roots of the equation.

The quadratic equation  $ax^2 + bx + c = 0$  with real coefficients has two roots given by  $\frac{-b + \sqrt{D}}{2a}$  and  $\frac{-b - \sqrt{D}}{2a}$ , where  $D = b^2 - 4ac$ , called the discriminant of the equation.

#### Notes

- When  $D = 0$ , roots of the quadratic equation are real and equal. When  $D > 0$ , roots are real and unequal. Further, if  $a, b, c \in \mathbf{Q}$  and  $D$  is a perfect square, then the roots of the equation are rational and unequal, and if  $a, b, c \in \mathbf{Q}$  and  $D$  is not a perfect square, then the roots are irrational and occur in pair.

When  $D < 0$ , roots of the quadratic equation are non real (or complex).

2. Let  $\alpha, \beta$  be the roots of the quadratic equation  $ax^2 + bx + c = 0$ , then sum of the roots

$$(\alpha + \beta) = \frac{-b}{a} \text{ and the product of the roots } (\alpha \cdot \beta) = \frac{c}{a}.$$

3. Let  $S$  and  $P$  be the sum of roots and product of roots, respectively, of a quadratic equation. Then the quadratic equation is given by  $x^2 - Sx + P = 0$ .

## 5.2 Solved Exmaples

### Short Answer Type

**Example 1** Evaluate :  $(1 + i)^6 + (1 - i)^3$

**Solution**  $(1 + i)^6 = \{(1 + i)^2\}^3 = (1 + i^2 + 2i)^3 = (1 - 1 + 2i)^3 = 8 i^3 = -8i$

and  $(1 - i)^3 = 1 - i^3 - 3i + 3i^2 = 1 + i - 3i - 3 = -2 - 2i$

Therefore,  $(1 + i)^6 + (1 - i)^3 = -8i - 2 - 2i = -2 - 10i$

**Example 2** If  $(x + iy)^{\frac{1}{3}} = a + ib$ , where  $x, y, a, b \in \mathbb{R}$ , show that  $\frac{x}{a} - \frac{y}{b} = -2(a^2 + b^2)$

**Solution**  $(x + iy)^{\frac{1}{3}} = a + ib$

$$\Rightarrow x + iy = (a + ib)^3$$

$$\begin{aligned} \text{i.e., } x + iy &= a^3 + i^3 b^3 + 3iab(a + ib) \\ &= a^3 - ib^3 + i3a^2b - 3ab^2 \\ &= a^3 - 3ab^2 + i(3a^2b - b^3) \end{aligned}$$

$$\Rightarrow x = a^3 - 3ab^2 \text{ and } y = 3a^2b - b^3$$

$$\text{Thus } \frac{x}{a} = a^2 - 3b^2 \text{ and } \frac{y}{b} = 3a^2 - b^2$$

$$\text{So, } \frac{x}{a} - \frac{y}{b} = a^2 - 3b^2 - 3a^2 + b^2 = -2a^2 - 2b^2 = -2(a^2 + b^2).$$

**Example 3** Solve the equation  $z^2 = \bar{z}$ , where  $z = x + iy$

**Solution**  $z^2 = \bar{z} \Rightarrow x^2 - y^2 + i2xy = x - iy$

$$\text{Therefore, } x^2 - y^2 = x \quad \dots (1) \quad \text{and} \quad 2xy = -y \quad \dots (2)$$

From (2), we have  $y = 0$  or  $x = -\frac{1}{2}$

When  $y = 0$ , from (1), we get  $x^2 - x = 0$ , i.e.,  $x = 0$  or  $x = 1$ .

When  $x = -\frac{1}{2}$ , from (1), we get  $y^2 = \frac{1}{4} + \frac{1}{2}$  or  $y^2 = \frac{3}{4}$ , i.e.,  $y = \pm \frac{\sqrt{3}}{2}$ .

Hence, the solutions of the given equation are

$$0 + i0, 1 + i0, -\frac{1}{2} + i\frac{\sqrt{3}}{2}, -\frac{1}{2} - i\frac{\sqrt{3}}{2}.$$

**Example 4** If the imaginary part of  $\frac{2z+1}{iz+1}$  is  $-2$ , then show that the locus of the point representing  $z$  in the argand plane is a straight line.

**Solution** Let  $z = x + iy$ . Then

$$\begin{aligned} \frac{2z+1}{iz+1} &= \frac{2(x+iy)+1}{i(x+iy)+1} = \frac{(2x+1)+i2y}{(1-y)+ix} \\ &= \frac{\{(2x+1)+i2y\}}{\{(1-y)+ix\}} \times \frac{\{(1-y)-ix\}}{\{(1-y)-ix\}} \\ &= \frac{(2x+1-y)+i(2y-2y^2-2x^2-x)}{1+y^2-2y+x^2} \end{aligned}$$

Thus 
$$\operatorname{Im}\left(\frac{2z+1}{iz+1}\right) = \frac{2y-2y^2-2x^2-x}{1+y^2-2y+x^2}$$

But 
$$\operatorname{Im}\left(\frac{2z+1}{iz+1}\right) = -2 \quad (\text{Given})$$

So 
$$\frac{2y-2y^2-2x^2-x}{1+y^2-2y+x^2} = -2$$

$\Rightarrow 2y - 2y^2 - 2x^2 - x = -2 - 2y^2 + 4y - 2x^2$   
i.e.,  $x + 2y - 2 = 0$ , which is the equation of a line.

**Example 5** If  $|z^2 - 1| = |z|^2 + 1$ , then show that  $z$  lies on imaginary axis.

**Solution** Let  $z = x + iy$ . Then  $|z^2 - 1| = |z|^2 + 1$

$$\begin{aligned} \Rightarrow & \quad |x^2 - y^2 - 1 + i2xy| = |x + iy|^2 + 1 \\ \Rightarrow & \quad (x^2 - y^2 - 1)^2 + 4x^2y^2 = (x^2 + y^2 + 1)^2 \\ \Rightarrow & \quad 4x^2 = 0 \quad \text{i.e.,} \quad x = 0 \end{aligned}$$

Hence  $z$  lies on  $y$ -axis.

**Example 6** Let  $z_1$  and  $z_2$  be two complex numbers such that  $\bar{z}_1 + i\bar{z}_2 = 0$  and  $\arg(z_1 z_2) = \pi$ . Then find  $\arg(z_1)$ .

**Solution** Given that  $\bar{z}_1 + i\bar{z}_2 = 0$

$$\begin{aligned} \Rightarrow & \quad z_1 = iz_2, \text{ i.e., } z_2 = -iz_1 \\ \text{Thus} & \quad \arg(z_1 z_2) = \arg z_1 + \arg(-iz_1) = \pi \\ \Rightarrow & \quad \arg(-iz_1^2) = \pi \\ \Rightarrow & \quad \arg(-i) + \arg(z_1^2) = \pi \\ \Rightarrow & \quad \arg(-i) + 2\arg(z_1) = \pi \\ \Rightarrow & \quad \frac{-\pi}{2} + 2\arg(z_1) = \pi \\ \Rightarrow & \quad \arg(z_1) = \frac{3\pi}{4} \end{aligned}$$

**Example 7** Let  $z_1$  and  $z_2$  be two complex numbers such that  $|z_1 + z_2| = |z_1| + |z_2|$ .

Then show that  $\arg(z_1) - \arg(z_2) = 0$ .

**Solution** Let  $z_1 = r_1(\cos\theta_1 + i\sin\theta_1)$  and  $z_2 = r_2(\cos\theta_2 + i\sin\theta_2)$

where  $r_1 = |z_1|$ ,  $\arg(z_1) = \theta_1$ ,  $r_2 = |z_2|$ ,  $\arg(z_2) = \theta_2$ .

$$\begin{aligned} \text{We have,} \quad |z_1 + z_2| &= |z_1| + |z_2| \\ &= |r_1(\cos\theta_1 + \cos\theta_2) + r_2(\cos\theta_2 + \sin\theta_2)| = r_1 + r_2 \\ &= r_1^2 + r_2^2 + 2r_1r_2\cos(\theta_1 - \theta_2) = (r_1 + r_2)^2 \Rightarrow \cos(\theta_1 - \theta_2) = 1 \\ &\Rightarrow \theta_1 - \theta_2 \text{ i.e. } \arg z_1 = \arg z_2 \end{aligned}$$

**Example 8** If  $z_1, z_2, z_3$  are complex numbers such that

$$|z_1| = |z_2| = |z_3| = \left| \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} \right| = 1, \text{ then find the value of } |z_1 + z_2 + z_3|.$$

**Solution**  $|z_1| = |z_2| = |z_3| = 1$



$$\Rightarrow |z_1|^2 = |z_2|^2 = |z_3|^2 = 1$$

$$\Rightarrow z_1 \bar{z}_1 = z_2 \bar{z}_2 = z_3 \bar{z}_3 = 1$$

$$\Rightarrow \bar{z}_1 = \frac{1}{z_1}, \bar{z}_2 = \frac{1}{z_2}, \bar{z}_3 = \frac{1}{z_3}$$

Given that  $\left| \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} \right| = 1$

$$\Rightarrow |\bar{z}_1 + \bar{z}_2 + \bar{z}_3| = 1, \text{ i.e., } |\overline{z_1 + z_2 + z_3}| = 1$$

$$\Rightarrow |z_1 + z_2 + z_3| = 1$$

**Example 9** If a complex number  $z$  lies in the interior or on the boundary of a circle of radius 3 units and centre  $(-4, 0)$ , find the greatest and least values of  $|z+1|$ .

**Solution** Distance of the point representing  $z$  from the centre of the circle is  $|z - (-4 + i0)| = |z+4|$ .

According to given condition  $|z+4| \leq 3$ .

$$\text{Now } |z+1| = |z+4-3| \leq |z+4| + |-3| \leq 3+3=6$$

Therefore, greatest value of  $|z+1|$  is 6.

Since least value of the modulus of a complex number is zero, the least value of  $|z+1|=0$ .

**Example 10** Locate the points for which  $3 < |z| < 4$

**Solution**  $|z| < 4 \Rightarrow x^2 + y^2 < 16$  which is the interior of circle with centre at origin and radius 4 units, and  $|z| > 3 \Rightarrow x^2 + y^2 > 9$  which is exterior of circle with centre at origin and radius 3 units. Hence  $3 < |z| < 4$  is the portion between two circles  $x^2 + y^2 = 9$  and  $x^2 + y^2 = 16$ .

**Example 11** Find the value of  $2x^4 + 5x^3 + 7x^2 - x + 41$ , when  $x = -2 - \sqrt{3}i$

**Solution**  $x + 2 = -\sqrt{3}i \Rightarrow x^2 + 4x + 7 = 0$

Therefore  $2x^4 + 5x^3 + 7x^2 - x + 41 = (x^2 + 4x + 7)(2x^2 - 3x + 5) + 6$   
 $= 0 \times (2x^2 - 3x + 5) + 6 = 6.$

**Example 12** Find the value of P such that the difference of the roots of the equation  $x^2 - Px + 8 = 0$  is 2.

**Solution** Let  $\alpha, \beta$  be the roots of the equation  $x^2 - Px + 8 = 0$

Therefore  $\alpha + \beta = P$  and  $\alpha \cdot \beta = 8$ .

Now  $\alpha - \beta = \pm \sqrt{(\alpha + \beta)^2 - 4\alpha\beta}$

Therefore  $2 = \pm \sqrt{P^2 - 32}$

$\Rightarrow P^2 - 32 = 4$ , i.e.,  $P = \pm 6$ .

**Example 13** Find the value of  $a$  such that the sum of the squares of the roots of the equation  $x^2 - (a - 2)x - (a + 1) = 0$  is least.

**Solution** Let  $\alpha, \beta$  be the roots of the equation

Therefore,  $\alpha + \beta = a - 2$  and  $\alpha\beta = -(a + 1)$

Now 
$$\begin{aligned} \alpha^2 + \beta^2 &= (\alpha + \beta)^2 - 2\alpha\beta \\ &= (a - 2)^2 + 2(a + 1) \\ &= (a - 1)^2 + 5 \end{aligned}$$

Therefore,  $\alpha^2 + \beta^2$  will be minimum if  $(a - 1)^2 = 0$ , i.e.,  $a = 1$ .

### Long Answer Type

**Example 14** Find the value of  $k$  if for the complex numbers  $z_1$  and  $z_2$ ,

$$|1 - \bar{z}_1 z_2|^2 - |z_1 - z_2|^2 = k(1 - |z_1|^2)(1 - |z_2|^2)$$

**Solution**

$$\begin{aligned} \text{L.H.S.} &= |1 - \bar{z}_1 z_2|^2 - |z_1 - z_2|^2 \\ &= (1 - \bar{z}_1 z_2)(\overline{1 - \bar{z}_1 z_2}) - (z_1 - z_2)(\overline{z_1 - z_2}) \\ &= (1 - \bar{z}_1 z_2)(1 - z_1 \bar{z}_2) - (z_1 - z_2)(\bar{z}_1 - \bar{z}_2) \\ &= 1 + z_1 \bar{z}_1 z_2 \bar{z}_2 - z_1 \bar{z}_1 - z_2 \bar{z}_2 \\ &= 1 + |z_1|^2 \cdot |z_2|^2 - |z_1|^2 - |z_2|^2 \\ &= (1 - |z_1|^2)(1 - |z_2|^2) \end{aligned}$$

$$\text{R.H.S.} = k(1 - |z_1|^2)(1 - |z_2|^2)$$

$\Rightarrow k = 1$

Hence, equating LHS and RHS, we get  $k = 1$ .

**Example 15** If  $z_1$  and  $z_2$  both satisfy  $z + \bar{z} = 2|z-1|$   $\arg(z_1 - z_2) = \frac{\pi}{4}$ , then find  $\text{Im}(z_1 + z_2)$ .

**Solution** Let  $z = x + iy$ ,  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$ .

$$\text{Then } z + \bar{z} = 2|z-1|$$

$$\Rightarrow (x + iy) + (x - iy) = 2|x-1+iy|$$

$$\Rightarrow 2x = 1 + y^2 \quad \dots (1)$$

Since  $z_1$  and  $z_2$  both satisfy (1), we have

$$2x_1 = 1 + y_1^2 \dots \text{ and } 2x_2 = 1 + y_2^2$$

$$\Rightarrow 2(x_1 - x_2) = (y_1 + y_2)(y_1 - y_2)$$

$$\Rightarrow 2 = (y_1 + y_2) \left( \frac{y_1 - y_2}{x_1 - x_2} \right) \quad \dots (2)$$

$$\text{Again } z_1 - z_2 = (x_1 - x_2) + i(y_1 - y_2)$$

$$\text{Therefore, } \tan \theta = \frac{y_1 - y_2}{x_1 - x_2}, \text{ where } \theta = \arg(z_1 - z_2)$$

$$\Rightarrow \tan \frac{\pi}{4} = \frac{y_1 - y_2}{x_1 - x_2} \quad \left( \text{since } \theta = \frac{\pi}{4} \right)$$

$$\text{i.e., } 1 = \frac{y_1 - y_2}{x_1 - x_2}$$

From (2), we get  $2 = y_1 + y_2$ , i.e.,  $\text{Im}(z_1 + z_2) = 2$

### Objective Type Questions

**Example 16** Fill in the blanks:

- The real value of 'a' for which  $3i^3 - 2a^2 + (1-a)i + 5$  is real is \_\_\_\_\_.
- If  $|z| = 2$  and  $\arg(z) = \frac{\pi}{4}$ , then  $z =$  \_\_\_\_\_.
- The locus of  $z$  satisfying  $\arg(z) = \frac{\pi}{3}$  is \_\_\_\_\_.
- The value of  $(-\sqrt{-1})^{4n-3}$ , where  $n \in \mathbf{N}$ , is \_\_\_\_\_.

- (v) The conjugate of the complex number  $\frac{1-i}{1+i}$  is \_\_\_\_\_.
- (vi) If a complex number lies in the third quadrant, then its conjugate lies in the \_\_\_\_\_.
- (vii) If  $(2+i)(2+2i)(2+3i)\dots(2+ni) = x+iy$ , then  $5.8.13 \dots (4+n^2) =$  \_\_\_\_\_.

**Solution**

- (i)  $3i^3 - 2ai^2 + (1-a)i + 5 = -3i + 2a + 5 + (1-a)i$   
 $= 2a + 5 + (-a-2)i$ , which is real if  $-a-2=0$  i.e.  $a=-2$ .
- (ii)  $z = |z| \left( \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) = 2 \left( \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) = \sqrt{2}(1+i)$
- (iii) Let  $z = x+iy$ . Then its polar form is  $z = r(\cos \theta + i \sin \theta)$ , where  $\tan \theta = \frac{y}{x}$

$\theta$  is  $\arg(z)$ . Given that  $\theta = \frac{\pi}{3}$ . Thus.

$$\tan \frac{\pi}{3} = \frac{y}{x} \Rightarrow y = \sqrt{3}x, \text{ where } x > 0, y > 0.$$

Hence, locus of  $z$  is the part of  $y = \sqrt{3}x$  in the first quadrant except origin.

- (iv) Here  $(-\sqrt{-1})^{4n-3} = (-i)^{4n-3} = (-i)^{4n} (-i)^{-3} = \frac{1}{(-i)^3}$   
 $= \frac{1}{-i^3} = \frac{1}{i} = \frac{i}{i^2} = -i$
- (v)  $\frac{1-i}{1+i} = \frac{1-i}{1+i} \times \frac{1-i}{1-i} = \frac{1+i^2-2i}{1-i^2} = \frac{1-1-2i}{1+1} = -i$

Hence, conjugate of  $\frac{1-i}{1+i}$  is  $i$ .

- (vi) Conjugate of a complex number is the image of the complex number about the  $x$ -axis. Therefore, if a number lies in the third quadrant, then its image lies in the second quadrant.
- (vii) Given that  $(2+i)(2+2i)(2+3i)\dots(2+ni) = x+iy$  ... (1)  
 $\Rightarrow \overline{(2+i)} \overline{(2+2i)} \overline{(2+3i)} \dots \overline{(2+ni)} = \overline{(x+iy)} = (x-iy)$   
 i.e.,  $(2-i)(2-2i)(2-3i)\dots(2-ni) = x-iy$  ... (2)

Multiplying (1) and (2), we get 5.8.13 ...  $(4 + n^2) = x^2 + y^2$ .

**Example 17** State true or false for the following:

- (i) Multiplication of a non-zero complex number by  $i$  rotates it through a right angle in the anti-clockwise direction.
- (ii) The complex number  $\cos\theta + i\sin\theta$  can be zero for some  $\theta$ .
- (iii) If a complex number coincides with its conjugate, then the number must lie on imaginary axis.
- (iv) The argument of the complex number  $z = (1 + i\sqrt{3})(1 + i)(\cos\theta + i\sin\theta)$  is  $\frac{7\pi}{12} + \theta$
- (v) The points representing the complex number  $z$  for which  $|z+1| < |z-1|$  lies in the interior of a circle.
- (vi) If three complex numbers  $z_1, z_2$  and  $z_3$  are in A.P., then they lie on a circle in the complex plane.
- (vii) If  $n$  is a positive integer, then the value of  $i^n + (i)^{n+1} + (i)^{n+2} + (i)^{n+3}$  is 0.

**Solution**

- (i) True. Let  $z = 2 + 3i$  be complex number represented by OP. Then  $iz = -3 + 2i$ , represented by OQ, where if OP is rotated in the anticlockwise direction through a right angle, it coincides with OQ.
- (ii) False. Because  $\cos\theta + i\sin\theta = 0 \Rightarrow \cos\theta = 0$  and  $\sin\theta = 0$ . But there is no value of  $\theta$  for which  $\cos\theta$  and  $\sin\theta$  both are zero.
- (iii) False, because  $x + iy = x - iy \Rightarrow y = 0 \Rightarrow$  number lies on  $x$ -axis.
- (iv) True,  $\arg(z) = \arg(1 + i\sqrt{3}) + \arg(1 + i) + \arg(\cos\theta + i\sin\theta)$   
 $\frac{\pi}{3} + \frac{\pi}{4} + \theta = \frac{7\pi}{12} + \theta$
- (v) False, because  $|x+iy+1| < |x+iy-1|$   
 $\Rightarrow (x+1)^2 + y^2 < (x-1)^2 + y^2$  which gives  $4x < 0$ .
- (vi) False, because if  $z_1, z_2$  and  $z_3$  are in A.P., then  $z_2 = \frac{z_1 + z_3}{2} \Rightarrow z_2$  is the midpoint of  $z_1$  and  $z_3$ , which implies that the points  $z_1, z_2, z_3$  are collinear.
- (vii) True, because  $i^n + (i)^{n+1} + (i)^{n+2} + (i)^{n+3}$   
 $= i^n(1 + i + i^2 + i^3) = i^n(1 + i - 1 - i)$   
 $= i^n(0) = 0$

**Example 18** Match the statements of column A and B.

**Column A**

**Column B**

- |   |                                      |
|---|--------------------------------------|
| (a) The value of $1+i^2 + i^4 + i^6 + \dots + i^{20}$ is  | (i) purely imaginary complex number  |
| (b) The value of $i^{-1097}$ is   | (ii) purely real complex number      |
| (c) Conjugate of $1+i$ lies in  | (iii) second quadrant                |
| (d) $\frac{1+2i}{1-i}$ lies in  | (iv) Fourth quadrant                 |
| (e) If $a, b, c \in \mathbb{R}$ and $b^2 - 4ac < 0$ , then the roots of the equation $ax^2 + bx + c = 0$ are non real (complex) and           | (v) may not occur in conjugate pairs |
| (f) If $a, b, c \in \mathbb{R}$ and $b^2 - 4ac > 0$ , and $b^2 - 4ac$ is a perfect square, then the roots of the equation $ax^2 + bx + c = 0$ | (vi) may occur in conjugate pairs    |

**Solution**

- (a)  $\Leftrightarrow$  (ii), because  $1 + i^2 + i^4 + i^6 + \dots + i^{20}$   
 $= 1 - 1 + 1 - 1 + \dots + 1 = 1$  (which is purely a real complex number)
- (b)  $\Leftrightarrow$  (i), because  $i^{-1097} = \frac{1}{(i)^{1097}} = \frac{1}{i^{4 \times 274 + 1}} = \frac{1}{\{(i)^4\}^{274} (i)} = \frac{1}{i} = \frac{i}{i^2} = -i$   
 which is purely imaginary complex number.
- (c)  $\Leftrightarrow$  (iv), conjugate of  $1 + i$  is  $1 - i$ , which is represented by the point  $(1, -1)$  in the fourth quadrant.
- (d)  $\Leftrightarrow$  (iii), because  $\frac{1+2i}{1-i} = \frac{1+2i}{1-i} \times \frac{1+i}{1+i} = \frac{-1+3i}{2} = -\frac{1}{2} + \frac{3}{2}i$ , which is represented by the point  $\left(-\frac{1}{2}, \frac{3}{2}\right)$  in the second quadrant.
- (e)  $\Leftrightarrow$  (vi), If  $b^2 - 4ac < 0 = D < 0$ , i.e., square root of  $D$  is a imaginary number, therefore, roots are  $x = \frac{-b \pm \text{Imaginary Number}}{2a}$ , i.e., roots are in conjugate pairs.

(f)  $\Leftrightarrow$  (v), Consider the equation  $x^2 - (5 + \sqrt{2})x + 5\sqrt{2} = 0$ , where  $a = 1$ ,  
 $b = -(5 + \sqrt{2})$ ,  $c = 5\sqrt{2}$ , clearly  $a, b, c \in \mathbb{R}$ .

$$\text{Now } D = b^2 - 4ac = \{-(5 + \sqrt{2})\}^2 - 4 \cdot 1 \cdot 5\sqrt{2} = (5 - \sqrt{2})^2.$$

Therefore  $x = \frac{5 + \sqrt{2} \pm 5 - \sqrt{2}}{2} = 5, \sqrt{2}$  which do not form a conjugate pair.

**Example 19** What is the value of  $\frac{i^{4n+1} - i^{4n-1}}{2}$ ?

**Solution**  $i$ , because  $\frac{i^{4n+1} - i^{4n-1}}{2} = \frac{i^{4n}i - i^{4n}i^{-1}}{2}$

$$= \frac{i - \frac{1}{i}}{2} = \frac{i^2 - 1}{2i} = \frac{-2}{2i} = i$$

**Example 20** What is the smallest positive integer  $n$ , for which  $(1 + i)^{2n} = (1 - i)^{2n}$ ?

**Solution**  $n = 2$ , because  $(1 + i)^{2n} = (1 - i)^{2n} = \left(\frac{1+i}{1-i}\right)^{2n} = 1$

$\Rightarrow (i)^{2n} = 1$  which is possible if  $n = 2$  ( $\because i^4 = 1$ )

**Example 21** What is the reciprocal of  $3 + \sqrt{7}i$ ?

**Solution** Reciprocal of  $z = \frac{\bar{z}}{|z|^2}$

$$\text{Therefore, reciprocal of } 3 + \sqrt{7}i = \frac{3 - \sqrt{7}i}{16} = \frac{3}{16} - \frac{\sqrt{7}i}{16}$$

**Example 22** If  $z_1 = \sqrt{3} + i\sqrt{3}$  and  $z_2 = \sqrt{3} + i$ , then find the quadrant in which

$\left(\frac{z_1}{z_2}\right)$  lies.

**Solution**  $\frac{z_1}{z_2} = \frac{\sqrt{3} + i\sqrt{3}}{\sqrt{3} + i} = \left(\frac{3 + \sqrt{3}}{4}\right) + \left(\frac{3 - \sqrt{3}}{4}\right)i$

which is represented by a point in first quadrant.

**Example 23** What is the conjugate of  $\frac{\sqrt{5+12i} + \sqrt{5-12i}}{\sqrt{5+12i} - \sqrt{5-12i}}$ ?

**Solution** Let

$$\begin{aligned} z &= \frac{\sqrt{5+12i} + \sqrt{5-12i}}{\sqrt{5+12i} - \sqrt{5-12i}} \times \frac{\sqrt{5+12i} + \sqrt{5-12i}}{\sqrt{5+12i} + \sqrt{5-12i}} \\ &= \frac{5+12i+5-12i+2\sqrt{25+144}}{5+12i-5+12i} \\ &= \frac{3}{2i} = \frac{3i}{-2} = 0 - \frac{3}{2}i \end{aligned}$$

Therefore, the conjugate of  $z = 0 + \frac{3}{2}i$

**Example 24** What is the principal value of amplitude of  $1 - i$ ?

**Solution** Let  $\theta$  be the principle value of amplitude of  $1 - i$ . Since

$$\tan \theta = -1 \Rightarrow \tan \theta = \tan\left(-\frac{\pi}{4}\right) \Rightarrow \theta = -\frac{\pi}{4}$$

**Example 25** What is the polar form of the complex number  $(i^{25})^3$ ?

**Solution**  $z = (i^{25})^3 = (i)^{75} = i^{4 \times 18 + 3} = (i^4)^{18} (i)^3$   
 $= i^3 = -i = 0 - i$

Polar form of  $z = r(\cos \theta + i \sin \theta)$

$$\begin{aligned} &= 1 \left\{ \cos\left(-\frac{\pi}{2}\right) + i \sin\left(-\frac{\pi}{2}\right) \right\} \\ &= \cos \frac{\pi}{2} - i \sin \frac{\pi}{2} \end{aligned}$$

**Example 26** What is the locus of  $z$ , if amplitude of  $z - 2 - 3i$  is  $\frac{\pi}{4}$ ?

**Solution** Let  $z = x + iy$ . Then  $z - 2 - 3i = (x - 2) + i(y - 3)$

Let  $\theta$  be the amplitude of  $z - 2 - 3i$ . Then  $\tan \theta = \frac{y-3}{x-2}$

$$\Rightarrow \tan \frac{\pi}{4} = \frac{y-3}{x-2} \left( \text{since } \theta = \frac{\pi}{4} \right)$$



$$\Rightarrow 1 = \frac{y-3}{x-2} \text{ i.e. } x - y + 1 = 0$$

Hence, the locus of  $z$  is a straight line.

**Example 27** If  $1 - i$ , is a root of the equation  $x^2 + ax + b = 0$ , where  $a, b \in \mathbf{R}$ , then find the values of  $a$  and  $b$ .

**Solution** Sum of roots  $\frac{-a}{1} = (1 - i) + (1 + i) \Rightarrow a = -2$ .

(since non real complex roots occur in conjugate pairs)

Product of roots,  $\frac{b}{1} = (1 - i)(1 + i) \Rightarrow b = 2$

Choose the correct options out of given four options in each of the Examples from 28 to 33 (M.C.Q.).

**Example 28**  $1 + i^2 + i^4 + i^6 + \dots + i^{2n}$  is

- (A) positive (B) negative  
(C) 0 (D) can not be evaluated

**Solution** (D),  $1 + i^2 + i^4 + i^6 + \dots + i^{2n} = 1 - 1 + 1 - 1 + \dots (-1)^n$

which can not be evaluated unless  $n$  is known.

**Example 29** If the complex number  $z = x + iy$  satisfies the condition  $|z + 1| = 1$ , then  $z$  lies on

- (A)  $x$ -axis  
(B) circle with centre  $(1, 0)$  and radius 1  
(C) circle with centre  $(-1, 0)$  and radius 1  
(D)  $y$ -axis

**Solution** (C),  $|z + 1| = 1 \Rightarrow |(x + 1) + iy| = 1$

$$\Rightarrow (x + 1)^2 + y^2 = 1$$

which is a circle with centre  $(-1, 0)$  and radius 1.

**Example 30** The area of the triangle on the complex plane formed by the complex numbers  $z$ ,  $-iz$  and  $z + iz$  is:

- (A)  $|z|^2$  (B)  $|\bar{z}|^2$   
(C)  $\frac{|z|^2}{2}$  (D) none of these

**Solution** (C), Let  $z = x + iy$ . Then  $-iz = y - ix$ . Therefore,

$$z + iz = (x - y) + i(x + y)$$

$$\text{Required area of the triangle} = \frac{1}{2}(x^2 + y^2) = \frac{|z|^2}{2}$$

**Example 31** The equation  $|z + 1 - i| = |z - 1 + i|$  represents a

- (A) straight line (B) circle  
(C) parabola (D) hyperbola

**Solution** (A),  $|z + 1 - i| = |z - 1 + i|$

$$\Rightarrow |z - (-1 + i)| = |z - (1 - i)|$$

$\Rightarrow$  PA = PB, where A denotes the point  $(-1, 1)$ , B denotes the point  $(1, -1)$  and P denotes the point  $(x, y)$

$\Rightarrow$  z lies on the perpendicular bisector of the line joining A and B and perpendicular bisector is a straight line.

**Example 32** Number of solutions of the equation  $z^2 + |z|^2 = 0$  is

- (A) 1 (B) 2  
(C) 3 (D) infinitely many

**Solution** (D),  $z^2 + |z|^2 = 0, z \neq 0$

$$\Rightarrow x^2 - y^2 + i2xy + x^2 + y^2 = 0$$

$$\Rightarrow 2x^2 + i2xy = 0 \Rightarrow 2x(x + iy) = 0$$

$$\Rightarrow x = 0 \text{ or } x + iy = 0 \text{ (not possible)}$$

Therefore,  $x = 0$  and  $z \neq 0$

So y can have any real value. Hence infinitely many solutions.

**Example 33** The amplitude of  $\sin \frac{\pi}{5} + i(1 - \cos \frac{\pi}{5})$  is

- (A)  $\frac{2\pi}{5}$  (B)  $\frac{\pi}{5}$  (C)  $\frac{\pi}{15}$  (D)  $\frac{\pi}{10}$

**Solution** (D), Here  $r \cos \theta = \sin \left( \frac{\pi}{5} \right)$  and  $r \sin \theta = 1 - \cos \frac{\pi}{5}$

$$\text{Therefore, } \tan \theta = \frac{1 - \cos \frac{\pi}{5}}{\sin \frac{\pi}{5}} = \frac{2 \sin^2 \left( \frac{\pi}{10} \right)}{2 \sin \left( \frac{\pi}{10} \right) \cdot \cos \left( \frac{\pi}{10} \right)}$$

$$\Rightarrow \tan \theta = \tan \left( \frac{\pi}{10} \right) \text{ i.e., } \theta = \frac{\pi}{10}$$

### 5.3 EXERCISE

#### Short Answer Type

- For a positive integer  $n$ , find the value of  $(1 - i)^n \left( 1 - \frac{1}{i} \right)^n$
- Evaluate  $\sum_{n=1}^{13} (i^n + i^{n+1})$ , where  $n \in \mathbf{N}$ .
- If  $\left( \frac{1+i}{1-i} \right)^3 - \left( \frac{1-i}{1+i} \right)^3 = x + iy$ , then find  $(x, y)$ .
- If  $\frac{(1+i)^2}{2-i} = x + iy$ , then find the value of  $x + y$ .
- If  $\left( \frac{1-i}{1+i} \right)^{100} = a + ib$ , then find  $(a, b)$ .
- If  $a = \cos \theta + i \sin \theta$ , find the value of  $\frac{1+a}{1-a}$ .
- If  $(1+i)z = (1-i)\bar{z}$ , then show that  $z = -i\bar{z}$ .
- If  $z = x + iy$ , then show that  $z\bar{z} + 2(z + \bar{z}) + b = 0$ , where  $b \in \mathbf{R}$ , represents a circle.
- If the real part of  $\frac{\bar{z}+2}{z-1}$  is 4, then show that the locus of the point representing  $z$  in the complex plane is a circle.
- Show that the complex number  $z$ , satisfying the condition  $\arg \left( \frac{z-1}{z+1} \right) = \frac{\pi}{4}$  lies on a circle.
- Solve the equation  $|z| = z + 1 + 2i$ .

**Long Answer Type**

12. If  $|z+1| = z + 2(1+i)$ , then find  $z$ .
13. If  $\arg(z-1) = \arg(z+3i)$ , then find  $x-1 : y$  where  $z = x + iy$
14. Show that  $\left| \frac{z-2}{z-3} \right| = 2$  represents a circle. Find its centre and radius.
15. If  $\frac{z-1}{z+1}$  is a purely imaginary number ( $z \neq -1$ ), then find the value of  $|z|$ .
16.  $z_1$  and  $z_2$  are two complex numbers such that  $|z_1| = |z_2|$  and  $\arg(z_1) + \arg(z_2) = \pi$ , then show that  $z_1 = -\bar{z}_2$ .
17. If  $|z_1| = 1$  ( $z_1 \neq -1$ ) and  $z_2 = \frac{z_1-1}{z_1+1}$ , then show that the real part of  $z_2$  is zero.
18. If  $z_1, z_2$  and  $z_3, z_4$  are two pairs of conjugate complex numbers, then find  $\arg\left(\frac{z_1}{z_4}\right) + \arg\left(\frac{z_2}{z_3}\right)$ .
19. If  $|z_1| = |z_2| = \dots = |z_n| = 1$ , then show that  $|z_1 + z_2 + z_3 + \dots + z_n| = \left| \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} + \dots + \frac{1}{z_n} \right|$ .
20. If for complex numbers  $z_1$  and  $z_2$ ,  $\arg(z_1) - \arg(z_2) = 0$ , then show that  $|z_1 - z_2| = |z_1| - |z_2|$
21. Solve the system of equations  $\operatorname{Re}(z^2) = 0$ ,  $|z| = 2$ .
22. Find the complex number satisfying the equation  $z + \sqrt{2}|(z+1)| + i = 0$ .
23. Write the complex number  $z = \frac{1-i}{\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}}$  in polar form.
24. If  $z$  and  $w$  are two complex numbers such that  $|zw| = 1$  and  $\arg(z) - \arg(w) = \frac{\pi}{2}$ , then show that  $\bar{z}w = -i$ .

**Objective Type Questions**

25. Fill in the blanks of the following

- (i) For any two complex numbers  $z_1, z_2$  and any real numbers  $a, b$ ,  
 $|az_1 - bz_2|^2 + |bz_1 + az_2|^2 = \dots$
- (ii) The value of  $\sqrt{-25} \times \sqrt{-9}$  is .....
- (iii) The number  $\frac{(1-i)^3}{1-i^3}$  is equal to .....
- (iv) The sum of the series  $i + i^2 + i^3 + \dots$  upto 1000 terms is .....
- (v) Multiplicative inverse of  $1 + i$  is .....
- (vi) If  $z_1$  and  $z_2$  are complex numbers such that  $z_1 + z_2$  is a real number, then  $z_2 = \dots$
- (vii)  $\arg(z) + \arg(\bar{z})$  ( $\bar{z} \neq 0$ ) is .....
- (viii) If  $|z+4| \leq 3$ , then the greatest and least values of  $|z+1|$  are ..... and .....
- (ix) If  $\left| \frac{z-2}{z+2} \right| = \frac{\pi}{6}$ , then the locus of  $z$  is .....
- (x) If  $|z| = 4$  and  $\arg(z) = \frac{5\pi}{6}$ , then  $z = \dots$

26. State True or False for the following :

- (i) The order relation is defined on the set of complex numbers.
- (ii) Multiplication of a non zero complex number by  $-i$  rotates the point about origin through a right angle in the anti-clockwise direction.
- (iii) For any complex number  $z$  the minimum value of  $|z| + |z-1|$  is 1.
- (iv) The locus represented by  $|z-1| = |z-i|$  is a line perpendicular to the join of (1, 0) and (0, 1).
- (v) If  $z$  is a complex number such that  $z \neq 0$  and  $\operatorname{Re}(z) = 0$ , then  $\operatorname{Im}(z^2) = 0$ .
- (vi) The inequality  $|z-4| < |z-2|$  represents the region given by  $x > 3$ .

(vii) Let  $z_1$  and  $z_2$  be two complex numbers such that  $|z_1 + z_2| = |z_1| + |z_2|$ , then  $\arg(z_1 - z_2) = 0$ .

(viii) 2 is not a complex number.

27. Match the statements of Column A and Column B.

**Column A**

**Column B**

- |   |  |
|---|--|
| (a) The polar form of $i + \sqrt{3}$ is             | (i) Perpendicular bisector of segment joining $(-2, 0)$ and $(2, 0)$     |
| (b) The amplitude of $-1 + \sqrt{-3}$ is            | (ii) On or outside the circle having centre at $(0, -4)$ and radius 3.   |
| (c) If $ z + 2  =  z - 2 $ , then locus of $z$ is   | (iii) $\frac{2\pi}{3}$   |
| (d) If $ z + 2i  =  z - 2i $ , then locus of $z$ is | (iv) Perpendicular bisector of segment joining $(0, -2)$ and $(0, 2)$ .  |
| (e) Region represented by $ z + 4i  \geq 3$ is      | (v) $2\left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}\right)$            |
| (f) Region represented by $ z + 4i  \leq 3$ is      | (vi) On or inside the circle having centre $(-4, 0)$ and radius 3 units. |
| (g) Conjugate of $\frac{1+2i}{1-i}$ lies in         | (vii) First quadrant   |
| (h) Reciprocal of $1 - i$ lies in                   | (viii) Third quadrant  |

28. What is the conjugate of  $\frac{2-i}{(1-2i)^2}$ ?

29. If  $|z_1| = |z_2|$ , is it necessary that  $z_1 = z_2$ ?

30. If  $\frac{(a^2+1)^2}{2a-i} = x + iy$ , what is the value of  $x^2 + y^2$ ?

31. Find  $z$  if  $|z| = 4$  and  $\arg(z) = \frac{5\pi}{6}$ .

32. Find  $\left| (1+i) \frac{(2+i)}{(3+i)} \right|$

33. Find principal argument of  $(1 + i\sqrt{3})^2$ .

34. Where does  $z$  lie, if  $\left| \frac{z-5i}{z+5i} \right| = 1$ .

Choose the correct answer from the given four options indicated against each of the Exercises from 35 to 50 (M.C.Q)

35.  $\sin x + i \cos 2x$  and  $\cos x - i \sin 2x$  are conjugate to each other for:

- (A)  $x = n\pi$  (B)  $x = \left(n + \frac{1}{2}\right) \frac{\pi}{2}$   
 (C)  $x = 0$  (D) No value of  $x$

36. The real value of  $\alpha$  for which the expression  $\frac{1-i\sin\alpha}{1+2i\sin\alpha}$  is purely real is :

- (A)  $(n+1) \frac{\pi}{2}$  (B)  $(2n+1) \frac{\pi}{2}$   
 (C)  $n\pi$  (D) None of these, where  $n \in \mathbb{N}$

37. If  $z = x + iy$  lies in the third quadrant, then  $\frac{\bar{z}}{z}$  also lies in the third quadrant if

- (A)  $x > y > 0$  (B)  $x < y < 0$   
 (C)  $y < x < 0$  (D)  $y > x > 0$

38. The value of  $(z+3)(\bar{z}+3)$  is equivalent to

- (A)  $|z+3|^2$  (B)  $|z-3|$   
 (C)  $z^2+3$  (D) None of these

39. If  $\left(\frac{1+i}{1-i}\right)^x = 1$ , then

- (A)  $x = 2n+1$  (B)  $x = 4n$   
 (C)  $x = 2n$  (D)  $x = 4n + 1$ , where  $n \in \mathbb{N}$

40. A real value of  $x$  satisfies the equation  $\left(\frac{3-4ix}{3+4ix}\right) = \alpha - i\beta$  ( $\alpha, \beta \in \mathbf{R}$ )  
if  $\alpha^2 + \beta^2 =$   
(A) 1 (B) -1 (C) 2 (D) -2
41. Which of the following is correct for any two complex numbers  $z_1$  and  $z_2$ ?  
(A)  $|z_1 z_2| = |z_1| |z_2|$  (B)  $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$   
(C)  $|z_1 + z_2| = |z_1| + |z_2|$  (D)  $|z_1 + z_2| \geq |z_1| - |z_2|$
42. The point represented by the complex number  $2 - i$  is rotated about origin through an angle  $\frac{\pi}{2}$  in the clockwise direction, the new position of point is:  
(A)  $1 + 2i$  (B)  $-1 - 2i$  (C)  $2 + i$  (D)  $-1 + 2i$
43. Let  $x, y \in \mathbf{R}$ , then  $x + iy$  is a non real complex number if:  
(A)  $x = 0$  (B)  $y = 0$  (C)  $x \neq 0$  (D)  $y \neq 0$
44. If  $a + ib = c + id$ , then  
(A)  $a^2 + c^2 = 0$  (B)  $b^2 + c^2 = 0$   
(C)  $b^2 + d^2 = 0$  (D)  $a^2 + b^2 = c^2 + d^2$
45. The complex number  $z$  which satisfies the condition  $\left|\frac{i+z}{i-z}\right| = 1$  lies on  
(A) circle  $x^2 + y^2 = 1$  (B) the  $x$ -axis  
(C) the  $y$ -axis (D) the line  $x + y = 1$ .
46. If  $z$  is a complex number, then  
(A)  $|z^2| > |z|^2$  (B)  $|z^2| = |z|^2$   
(C)  $|z^2| < |z|^2$  (D)  $|z^2| \geq |z|^2$
47.  $|z_1 + z_2| = |z_1| + |z_2|$  is possible if  
(A)  $z_2 = \bar{z}_1$  (B)  $z_2 = \frac{1}{z_1}$   
(C)  $\arg(z_1) = \arg(z_2)$  (D)  $|z_1| = |z_2|$



48. The real value of  $\theta$  for which the expression  $\frac{1+i\cos\theta}{1-2i\cos\theta}$  is a real number is:
- (A)  $n\pi + \frac{\pi}{4}$  (B)  $n\pi + (-1)^n \frac{\pi}{4}$   
(C)  $2n\pi \pm \frac{\pi}{2}$  (D) none of these.
49. The value of  $\arg(x)$  when  $x < 0$  is:
- (A) 0 (B)  $\frac{\pi}{2}$   
(C)  $\pi$  (D) none of these
50. If  $f(z) = \frac{7-z}{1-z^2}$ , where  $z = 1 + 2i$ , then  $|f(z)|$  is
- (A)  $\frac{|z|}{2}$  (B)  $|z|$   
(C)  $2|z|$  (D) none of these.

