

The p-Block Elements

I. Multiple Choice Questions (Type-I)

- **1.** On addition of conc. H₂SO₄ to a chloride salt, colourless fumes are evolved but in case of iodide salt, violet fumes come out. This is because
 - (i) H_2SO_4 reduces HI to I_2
 - (ii) HI is of violet colour
 - (iii) HI gets oxidised to I₂
 - (iv) HI changes to HIO₃
- **2.** In qualitative analysis when H₂S is passed through an aqueous solution of salt acidified with dil. HCl, a black precipitate is obtained. On boiling the precipitate with dil. HNO₃, it forms a solution of blue colour. Addition of excess of aqueous solution of ammonia to this solution gives ______.
 - (i) deep blue precipitate of Cu (OH)₂
 - (ii) deep blue solution of $[Cu (NH_3)_4]^{2+}$
 - (iii) deep blue solution of Cu(NO₃)₂
 - (iv) deep blue solution of Cu(OH)₂.Cu(NO₃)₂
- **3.** In a cyclotrimetaphosphoric acid molecule, how many single and double bonds are present?
 - (i) 3 double bonds; 9 single bonds
 - (ii) 6 double bonds; 6 single bonds
 - (iii) 3 double bonds; 12 single bonds
 - (iv) Zero double bonds; 12 single bonds
- **4.** Which of the following elements can be involved in $p\pi$ - $d\pi$ bonding?
 - (i) Carbon
 - (ii) Nitrogen

- (iii) Phosphorus
- (iv) Boron
- **5.** Which of the following pairs of ions are isoelectronic and isostructural?
 - (i) CO_3^{2-}, NO_3^{-}
 - (ii) ClO_3^- , CO_3^{2-}
 - (iii) SO_3^{2-} , NO_3^{-}
 - (iv) ClO_3^-, SO_3^2
- **6.** Affinity for hydrogen decreases in the group from fluorine to iodine. Which of the halogen acids should have highest bond dissociation enthalpy?
 - (i) HF
 - (ii) HCl
 - (iii) HBr
 - (iv) HI
- **7.** Bond dissociation enthalpy of E—H (E = element) bonds is given below. Which of the compounds will act as strongest reducing agent?

Compound	NH_3	$\mathbf{PH}_{_{3}}$	\mathbf{AsH}_3	SbH_3
$\Delta_{ m diss}$ (E—H)/kJ mol $^{-1}$	389	322	297	255

- (i) NH₃
- (ii) PH₃
- (iii) AsH₃
- (iv) SbH₃
- **8.** On heating with concentrated NaOH solution in an inert atmosphere of CO₂, white phosphorus gives a gas. Which of the following statement is **incorrect** about the gas?
 - (i) It is highly poisonous and has smell like rotten fish.
 - (ii) It's solution in water decomposes in the presence of light.
 - (iii) It is more basic than NH₃.
 - (iv) It is less basic than NH_3 .
- **9.** Which of the following acids forms three series of salts?
 - (i) H_3PO_9
 - (ii) H_oBO_o
 - (iii) H₃PO₄
 - (iv) H_3PO_3
- **10.** Strong reducing behaviour of H₃PO₂ is due to
 - (i) Low oxidation state of phosphorus
 - (ii) Presence of two -OH groups and one P-H bond

- (iii) Presence of one –OH group and two P–H bonds
- (iv) High electron gain enthalpy of phosphorus
- **11.** On heating lead nitrate forms oxides of nitrogen and lead. The oxides formed are _____.
 - (i) N₂O, PbO
 - (ii) NO₂, PbO
 - (iii) NO, PbO
 - (iv) NO, PbO₂
- **12.** Which of the following elements does not show allotropy?
 - (i) Nitrogen
 - (ii) Bismuth
 - (iii) Antimony
 - (iv) Arsenic
- 13. Maximum covalency of nitrogen is ______
 - (i) 3
 - (ii) 5
 - (iii) 4
 - (iv) 6
- **14.** Which of the following statements is wrong?
 - (i) Single N-N bond is stronger than the single P-P bond.
 - (ii) ${\rm PH_3}$ can act as a ligand in the formation of coordination compound with transition elements.
 - (iii) NO₂ is paramagnetic in nature.
 - (iv) Covalency of nitrogen in N_2O_5 is four.
- **15.** A brown ring is formed in the ring test for NO₃ ion. It is due to the formation of
 - (i) $[Fe(H_2O)_5 (NO)]^{2+}$
 - (ii) FeSO₄.NO₂
 - (iii) $[Fe(H_2O)_4(NO)_2]^{2+}$
 - (iv) FeSO₄.HNO₃
- **16.** Elements of group-15 form compounds in +5 oxidation state. However, bismuth forms only one well characterised compound in +5 oxidation state. The compound is
 - (i) Bi_2O_5
 - (ii) BiF₅
 - (iii) BiCl₅
 - (iv) $Bi_{2}S_{5}$

- 17. On heating ammonium dichromate and barium azide separately we get (i) N_o in both cases N_{2} with ammonium dichromate and NO with barium azide N₂O with ammonium dichromate and N₂ with barium azide N₂O with ammonium dichromate and NO₂ with barium azide In the preparation of HNO₃, we get NO gas by catalytic oxidation of ammonia. The moles of NO produced by the oxidation of two moles of NH₃ will be _____. (i) (ii) 3 4 (iii) (iv) 6 **19.** The oxidation state of central atom in the anion of compound NaH₂PO₂ will (i) +3 (ii) +5 (iii) +1(iv) -3**20.** Which of the following is **not** tetrahedral in shape? (i) NH_{A}^{\dagger} (ii) SiCl, SF_4 (iii) SO, 2-(iv) **21.** Which of the following are peroxoacids of sulphur? H₂SO₅ and H₂S₂O₈ (ii) H_2SO_5 and $H_2S_2O_7$ (iii) $H_2S_2O_7$ and $H_2S_2O_8$ $H_2S_2O_6$ and $H_2S_2O_7$ Hot conc. H₂SO₄ acts as moderately strong oxidising agent. It oxidises both metals and nonmetals. Which of the following element is oxidised by conc. H₂SO₄ into two gaseous products? (i) Cu S (ii) (iii) C (iv) Zn A black compound of manganese reacts with a halogen acid to give greenish yellow gas. When excess of this gas reacts with NH3 an unstable trihalide is formed. In this process the oxidation state of nitrogen changes from _ - 3 to +3 (i)
 - 93 p-Block Elements

(ii)

(iii) (iv) -3 to 0-3 to +5

0 to -3

- **24.** In the preparation of compounds of Xe, Bartlett had taken O_2^+ Pt F_6^- as a base compound. This is because
 - (i) both O₂ and Xe have same size.
 - (ii) both O_2 and Xe have same electron gain enthalpy.
 - (iii) both O₂ and Xe have almost same ionisation enthalpy.
 - (iv) both Xe and O₂ are gases.
- **25.** In solid state PCl₅ is a _____.
 - (i) covalent solid
 - (ii) octahedral structure
 - (iii) ionic solid with [PCl₆]⁺octahedral and [PCl₄]⁻tetrahedra
 - (iv) ionic solid with [PCl₄] tetrahedral and [PCl₆] octahedra
- **26.** Reduction potentials of some ions are given below. Arrange them in decreasing order of oxidising power.

Ion $ClO_4^ IO_4^ BrO_4^-$ **Reduction** $E^{\ominus}=1.19V$ $E^{\ominus}=1.65V$ $E^{\ominus}=1.74V$ **potential** E^{\ominus}/V

- (i) $ClO_4^- > IO_4^- > BrO_4^-$
- (ii) $IO_4^- > BrO_4^- > ClO_4^-$
- (iii) $BrO_4^- > IO_4^- > ClO_4^-$
- (iv) $BrO_4^- > ClO_4^- > IO_4^-$
- **27.** Which of the following is isoelectronic pair?
 - (i) ICl₂, ClO₂
 - (ii) BrO_2^- , BrF_2^+
 - (iii) ClO₂, BrF
 - (iv) CN^{-} , O_3

II. Multiple Choice Questions (Type-II)

Note: In the following questions two or more options may be correct.

- **28.** If chlorine gas is passed through hot NaOH solution, two changes are observed in the oxidation number of chlorine during the reaction. These are _____ and _____.
 - (i) 0 to +5
 - (ii) 0 to +3
 - (iii) 0 to -1
 - (iv) 0 to +1

- **29.** Which of the following options are **not** in accordance with the property mentioned against them?
 - (i) $F_2 > Cl_2 > Br_2 > I_2$ Oxidising power.
 - (ii) MI > MBr > MCl > MF Ionic character of metal halide.
 - (iii) $F_2 > Cl_2 > Br_2 > I_2$ Bond dissociation enthalpy.
 - (iv) HI < HBr < HCl < HF Hydrogen-halogen bond strength.
- **30.** Which of the following is correct for P_4 molecule of white phosphorus?
 - (i) It has 6 lone pairs of electrons.
 - (ii) It has six P-P single bonds.
 - (iii) It has three P-P single bonds.
 - (iv) It has four lone pairs of electrons.
- **31.** Which of the following statements are correct?
 - (i) Among halogens, radius ratio between iodine and fluorine is maximum.
 - (ii) Leaving F—F bond, all halogens have weaker X—X bond than X—X' bond in interhalogens.
 - (iii) Among interhalogen compounds maximum number of atoms are present in iodine fluoride.
 - (iv) Interhalogen compounds are more reactive than halogen compounds.
- **32.** Which of the following statements are correct for SO₂ gas?
 - (i) It acts as bleaching agent in moist conditions.
 - (ii) It's molecule has linear geometry.
 - (iii) It's dilute solution is used as disinfectant.
 - (iv) It can be prepared by the reaction of dilute H_2SO_4 with metal sulphide.
- **33.** Which of the following statements are correct?
 - (i) All the three N—O bond lengths in HNO₃ are equal.
 - (ii) All P—Cl bond lengths in PCl_5 molecule in gaseous state are equal.
 - (iii) P_4 molecule in white phohsphorus have angular strain therefore white phosphorus is very reactive.
 - (iv) PCl is ionic in solid state in which cation is tetrahedral and anion is octahedral.
- **34.** Which of the following orders are correct as per the properties mentioned against each?
 - (i) $As_2O_3 < SiO_2 < P_2O_3 < SO_2$ Acid strength.
 - (ii) $AsH_3 < PH_3 < NH_3$ Enthalpy of vapourisation.
 - (iii) S < O < Cl < F More negative electron gain enthalpy.
 - (iv) $H_2O > H_2S > H_2Se > H_2Te$ Thermal stability.

- **35.** Which of the following statements are correct?
 - (i) S–S bond is present in $H_2S_2O_6$.
 - (ii) In peroxosulphuric acid (H₂SO₅) sulphur is in +6 oxidation state.
 - (iii) Iron powder along with Al_2O_3 and K_2O is used as a catalyst in the preparation of NH_3 by Haber's process.
 - (iv) Change in enthalpy is positive for the preparation of SO_3 by catalytic oxidation of SO_2 .
- **36.** In which of the following reactions conc. H₂SO₄ is used as an oxidising reagent?
 - (i) $CaF_2 + H_2SO_4 \longrightarrow CaSO_4 + 2HF$
 - (ii) $2HI + H_2SO_4 \longrightarrow I_2 + SO_2 + 2H_2O$
 - (iii) $Cu + 2H_2SO_4 \longrightarrow CuSO_4 + SO_2 + 2H_2O$
 - (iv) $NaCl + H_2SO_4 \longrightarrow NaHSO_4 + HCl$
- **37.** Which of the following statements are true?
 - (i) Only type of interactions between particles of noble gases are due to weak dispersion forces.
 - (ii) Ionisation enthalpy of molecular oxygen is very close to that of xenon.
 - (iii) Hydrolysis of XeF₆ is a redox reaction.
 - (iv) Xenon fluorides are not reactive.

III. Short Answer Type

- **38.** In the preparation of H₂SO₄ by Contact Process, why is SO₃ not absorbed directly in water to form H₂SO₄?
- **39.** Write a balanced chemical equation for the reaction showing catalytic oxidation of NH₃ by atmospheric oxygen.
- **40.** Write the structure of pyrophosphoric acid.
- **41.** PH_3 forms bubbles when passed slowly in water but NH_3 dissolves. Explain why?
- **42.** In PCl_5 , phosphorus is in sp^3d hybridised state but all its five bonds are not equivalent. Justify your answer with reason.
- **43.** Why is nitric oxide paramagnetic in gaseous state but the solid obtained on cooling it is diamagnetic?
- **44.** Give reason to explain why ClF₃ exists but FCl₃ does not exist.
- **45.** Out of H₂O and H₂S, which one has higher bond angle and why?
- **46.** SF_6 is known but SCl_6 is not. Why?
- **47.** On reaction with Cl₂, phosphorus forms two types of halides 'A' and 'B'. Halide A is yellowish-white powder but halide 'B' is colourless oily liquid. Identify A and B and write the formulas of their hydrolysis products.

- **48.** In the ring test of NO_3^- ion, Fe^{2+} ion reduces nitrate ion to nitric oxide, which combines with Fe^{2+} (aq) ion to form brown complex. Write the reactions involved in the formation of brown ring.
- **49.** Explain why the stability of oxoacids of chlorine increases in the order given below:

HClO < HClO₂ < HClO₃ < HClO₄

- **50.** Explain why ozone is thermodynamically less stable than oxygen.
- **51.** P_4O_6 reacts with water according to equation $P_4O_6 + 6H_2O \longrightarrow 4H_3PO_3$. Calculate the volume of 0.1 M NaOH solution required to neutralise the acid formed by dissolving 1.1 g of P_4O_6 in H_2O .
- **52.** White phosphorus reacts with chlorine and the product hydrolyses in the presence of water. Calculate the mass of HCl obtained by the hydrolysis of the product formed by the reaction of 62 g of white phosphorus with chlorine in the presence of water.
- **53.** Name three oxoacids of nitrogen. Write the disproportionation reaction of that oxoacid of nitrogen in which nitrogen is in +3 oxidation state.
- **54.** Nitric acid forms an oxide of nitrogen on reaction with P_4O_{10} . Write the reaction involved. Also write the resonating structures of the oxide of nitrogen formed.
- **55.** Phosphorus has three allotropic forms (i) white phosphorus (ii) red phosphorus and (iii) black phosphorus. Write the difference between white and red phosphorus on the basis of their structure and reactivity.
- **56.** Give an example to show the effect of concentration of nitric acid on the formation of oxidation product.
- **57.** PCl_5 reacts with finely divided silver on heating and a white silver salt is obtained, which dissolves on adding excess aqueous NH_3 solution. Write the reactions involved to explain what happens.
- **58.** Phosphorus forms a number of oxoacids. Out of these oxoacids phosphinic acid has strong reducing property. Write its structure and also write a reaction showing its reducing behaviour.

IV. Matching Type

Note: Match the items of Column I and Column II in the following questions.

59. Match the compounds given in Column I with the hybridisation and shape given in Column II and mark the correct option.

Column I

- (A) $\operatorname{Xe} F_6$
- (B) Xe O₃
- (C) $Xe OF_4$
- (D) Xe F₄

Column II

- (1) sp^3d^3 distorted octahedral
- (2) sp^3d^2 square planar
- (3) sp^3 pyramidal
- (4) sp³ d² square pyramidal

Code:

(i)	A (1)	B (3)	C (4)	D (2)
(ii)	A (1)	B (2)	C (4)	D (3)
(iii)	A (4)	B (3)	C (1)	D (2)
(iv)	A (4)	B (1)	C (2)	D (3)

60. Match the formulas of oxides given in Column I with the type of oxide given in Column II and mark the correct option.

Column I **Column II** (A) Pb_3O_4 (1) Neutral oxide (B) N₂O (2)Acidic oxide (C) $Mn_{2}O_{7}$ (3)Basic oxide (D) Bi_2O_3 (4)Mixed oxide Code: (i) A (1) B (2) C (3) D (4) (ii) A (4) B (1) C (2) D (3) A (3) (iii) B (2) C (4) D (1) (iv) A (4) B (3) C(1) D (2)

61. Match the items of Columns I and II and mark the correct option.

Column I

- (A) H_2SO_4
- (B) CCl₃NO₂
- (C) Cl₂
- (D) Sulphur

Column II

- (1) Highest electron gain enthalpy
- (2) Chalcogen
- (3) Tear gas
- (4) Storage batteries

Code:

(i)	A (4)	B (3)	C (1)	D (2)
(ii)	A (3)	B (4)	C (1)	D (2)
(iii)	A (4)	B(1)	C (2)	D (3)
(iv)	A (2)	B (1)	C (3)	D (4)

62. Match the species given in Column I with the shape given in Column II and mark the correct option.

Column I

- (A) SF₄
- (B) BrF₃
- (C) BrO_3^-
- (D) NH₄

Column II

- (1) Tetrahedral
- (2) Pyramidal
- (3) Sea-saw shaped
- (4) Bent T-shaped

Code:

(i)	A (3)	B (2)	C (1)	D (4)
(ii)	A (3)	B (4)	C (2)	D (1)
(iii)	A (1)	B (2)	C (3)	D (4)
(iv)	A (1)	B (4)	C (3)	D (2)

63. Match the items of Columns I and II and mark the correct option.

Column I Column II

- (A) Its partial hydrolysis does not (1) He change oxidation state of central atom
- (B) It is used in modern diving apparatus (2) XeF₆
- (C) It is used to provide inert atmosphere $\qquad \qquad \text{(3)} \quad \text{XeF}_4$ for filling electrical bulbs
- (D) Its central atom is in sp^3d^2 hybridisation (4) Ar

Code:

(i)	A (1)	B (4)	C (2)	D (3)
(ii)	A (1)	B (2)	C (3)	D (4)
(iii)	A (2)	B(1)	C (4)	D (3)
(iv)	A (1)	B (3)	C(2)	D (4)

V. Assertion and Reason Type

Note: In the following questions a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.

- (i) Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.
- (ii) Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.
- (iii) Assertion is correct, but reason is wrong statement.
- (iv) Assertion is wrong but reason is correct statement.
- (v) Both assertion and reason are wrong statements.
- **64. Assertion** : N_2 is less reactive than P_4 .
 - **Reason**: Nitrogen has more electron gain enthalpy than phosphorus.
- **65. Assertion** : HNO₃ makes iron passive.
 - **Reason**: HNO₃ forms a protective layer of ferric nitrate on the surface of iron.

66. Assertion : HI cannot be prepared by the reaction of KI with concentrated

H₂SO₄

Reason: HI has lowest H–*X* bond strength among halogen acids.

67. Assertion : Both rhombic and monoclinic sulphur exist as S_8 but oxygen

exists as O_2 .

Reason : Oxygen forms $p\pi - p\pi$ multiple bond due to small size and

small bond length but $p\pi$ – $p\pi$ bonding is not possible in

sulphur.

68. Assertion : NaCl reacts with concentrated H₂SO₄ to give colourless fumes

with pungent smell. But on adding MnO₂ the fumes become

greenish yellow.

Reason: MnO₂ oxidises HCl to chlorine gas which is greenish yellow.

69. Assertion : SF₆ cannot be hydrolysed but SF₄ can be.

Reason: Six F atoms in SF₆ prevent the attack of H₂O on sulphur

atom of SF₆.

VI. Long Answer Type

70. An amorphous solid "A" burns in air to form a gas "B" which turns lime water milky. The gas is also produced as a by-product during roasting of sulphide ore. This gas decolourises acidified aqueous ${\rm KMnO_4}$ solution and reduces ${\rm Fe^{3+}}$ to ${\rm Fe^{2+}}$. Identify the solid "A" and the gas "B" and write the reactions involved.

- **71.** On heating lead (II) nitrate gives a brown gas "A". The gas "A" on cooling changes to colourless solid "B". Solid "B" on heating with NO changes to a blue solid 'C'. Identify 'A', 'B' and 'C' and also write reactions involved and draw the structures of 'B' and 'C'.
- **72.** On heating compound (A) gives a gas (B) which is a constituent of air. This gas when treated with 3 mol of hydrogen (H₂) in the presence of a catalyst gives another gas (C) which is basic in nature. Gas C on further oxidation in moist condition gives a compound (D) which is a part of acid rain. Identify compounds (A) to (D) and also give necessary equations of all the steps involved.

ANSWERS

I. Multiple Choice Questions (Type-I)

1. (iii)	2. (ii)	3. (i)	4. (iii)	5. (i)	6. (i)
7. (iv)	8. (iii)	9. (iii)	10. (iii)	11. (ii)	12. (i)
13. (iii)	14. (i)	15. (i)	16. (ii)	17. (i)	18. (i)
19. (iii)	20. (iii)	21. (i)	22. (iii)	23. (i)	24. (iii)
25. (iv)	26. (iii)	27. (ii)			

II. Multiple Choice Questions (Type-II)

28. (i), (iii)	29. (ii), (iii)	30. (ii), (iv)	31. (i), (iii), (iv)
32. (i), (iii)	33. (iii), (iv)	34. (i), (iv)	35. (i), (ii)
36. (ii), (iii)	37. (i), (ii)		

III. Short Answer Type

38. Acid fog is formed, which is difficult to condense.

39.
$$4\text{NH}_3 + 5\text{O}_2 \xrightarrow{\text{Pt/Rh gauge catalyst}} 4\text{NO} + 6\text{H}_2\text{O}$$
(From air)

- 41. NH_3 forms hydrogen bonds with water therefore it is soluble in it but PH_3 cannot form hydrogen bond with water so it escapes as gas.
- 42. **[Hint:** It has trigonal bipyramidal geometry]
- 43. In gaseous state $\mathrm{NO_2}$ exists as monomer which has one unpaired electron but in solid state it dimerises to $\mathrm{N_2O_4}$ so no unpaired electron is left hence solid form is diamagnetic.
- 44. Because fluorine is more electronegative as compared to chlorine.
- 45. Bond angle of $\rm H_2O$ is larger, because oxygen is more electronegative than sulphur therefore bond pair electron of O–H bond will be closer to oxygen and there will be more bond-pair bond-pair repulsion between bond pairs of two O–H bonds.
- 46. Due to small size of fluorine $\sin F$ ion can be accommodated around sulphur whereas chloride ion is comparatively larger in size, therefore, there will be interionic repulsion.

47. A is PCl₅ (It is yellowish white powder)

$$P_4 + 10Cl_2 \longrightarrow 4PCl_5$$

B is PCl₃ (It is a colourless oily liquid)

$$P_4 + 6Cl_2 \longrightarrow 4PCl_3$$

Hydrolysis products are formed as follows:

$$PCl_3 + 3H_2O \longrightarrow H_3PO_3 + 3HCl$$

$$PCl_5 + 4H_9O \longrightarrow H_3PO_4 + 5HCl$$

48. $NO_3^- + 3Fe^{2+} + 4H^+ \longrightarrow NO + 3Fe^{3+} + 2H_2O$

$$[\mathrm{Fe}(\mathrm{H_2O)_6}]^{2^+} + \mathrm{NO} \longrightarrow [\mathrm{Fe}(\mathrm{H_2O)_5}(\mathrm{NO})]^{2^+} + \mathrm{H_2O}$$

(brown complex)

49. Oxygen is more electronegative than chlorine, therefore dispersal of negative charge present on chlorine increases from ${\rm ClO}^{\scriptscriptstyle -}$ to ${\rm ClO}_4^{\scriptscriptstyle -}$ ion because number of oxygen atoms attached to chlorine is increasing. Therefore, stability of ions will increase in the order given below :

$$\text{ClO}^{-} < \text{ClO}_{2}^{-} < \text{ClO}_{3}^{-} < \text{ClO}_{4}^{-}$$

Thus due to increase in stability of conjugate base, acidic strength of corresponding acid increases in the following order

$$\mathrm{HClO} < \mathrm{HClO}_2 < \mathrm{HClO}_3 < \mathrm{HClO}_4$$

- 50. See the NCERT textbook for Class XII, page 186.
- 51. $P_4O_6 + 6H_2O \longrightarrow 4H_2PO_3$

$$H_3PO_3 + 2NaOH \longrightarrow Na_2 HPO_3 + 2H_2O] \times 4$$
 (Neutralisation reaction)

$$P_4O_6 + 8NaOH \longrightarrow 4Na_2 HPO_4 + 2H_2O$$

1 mol 8 mol

Product formed by 1 mol of P₄O₆ is neutralised by 8 mols of NaOH

∴ Product formed by $\frac{1.1}{220}$ mol of P_4O_6 will be neutralised by $\frac{1.1}{220} \times 8$ mol

of NaOH

Molarity of NaOH solution is 0.1M

 \Rightarrow 0.1 mol NaOH is present in 1 L solution

∴
$$\frac{1.1}{220} \times 8$$
 mol NaOH is present in $\frac{1.1 \times 8}{220 \times 0.1}$ L = $\frac{88}{220}$ L = $\frac{4}{10}$ L = 0.4 L =

400 mL of NaOH solution.

52. $P_4 + 6Cl_2 \longrightarrow 4PCl_3$ $PCl_3 + 3H_2O \longrightarrow H_3PO_3 + 3HCl] \times 4$ $P_4 + 6Cl_2 + 12H_2O \longrightarrow 4H_3PO_3 + 12HCl$

1 mol of white phosphorus produces 12 mol of HCl

62g of white phosphorus has been taken which is equivalent to $\frac{62}{124} = \frac{1}{2}$ mol.

Therefore 6 mol HCl will be formed.

Mass of 6 mol HCl = $6 \times 36.5 = 219.0$ g HCl

- 53. Three oxoacids of nitrogen are
 - (i) HNO₂, Nitrous acid
 - (ii) HNO₃, Nitric acid
 - (iii) Hyponitrous acid, H₂N₂O₂

$$3 \text{HNO}_2 \xrightarrow{\quad \text{Disproportionation} \quad} \text{HNO}_3 + \text{H}_2\text{O} + 2 \text{NO}$$

54.
$$4HNO_3 + P_4O_{10} \longrightarrow 4HPO_3 + 2N_2O_5$$

- 55. (a) Structures (See NCERT textbook for Class XII)
 - White phosphorus is discrete tetrahedral molecule. Thus it has tetrahedral structure with six P–P bonds.
 - Red phosphorus has polymeric structure in which P_4 tetrahedra are linked together through P—P bonds to form chain.
 - (b) Reactivity

White phosphorus is much more reactive than red phosphorus. This is because in white phosphorus there is angular strain in P_4 molecules because the bond angles are only of 60° .

56. Dilute and concentrated nitric acid give different oxidation products on reaction with copper metal.

$$3\text{Cu} + 8\text{HNO}_3 \text{ (dil.)} \longrightarrow 3\text{Cu(NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O}$$

 $\text{Cu} + 4\text{HNO}_3 \text{ (Conc.)} \longrightarrow 3\text{Cu(NO}_3)_2 + 2\text{NO} + 2\text{H}_2\text{O}$

57.
$$PCl_5 + 2Ag \longrightarrow 2AgCl + PCl_3$$

$$AgCl + 2NH_3(aq) \longrightarrow [Ag(NH_3)_2]^{\dagger}Cl$$

(soluble complex)

 $58. \hspace{0.1in}$ Structure of phosphinic acid (Hypophosphorous acid) is as follows :

Reducing behaviour of phosphinic acid is observable in the reaction with silver nitrate given below:

$$4 \mathrm{AgNO_3} + 2 \mathrm{H_2O} + \mathrm{H_3PO_2} {\longrightarrow} 4 \mathrm{Ag} + 4 \mathrm{HNO_3} + \mathrm{H_3PO_4}$$

IV. Matching Type

59. (i)

60. (ii)

61. (i)

62. (ii)

63. (iii)

V. Assertion and Reason Type

64. (iii)

65. (iii)

66. (ii)

67. (i)

68. (i)

69. (i)

VI. Long Answer Type

70. 'A' is
$$S_8$$

70. 'A' is S_8 'B' is SO_2 gas

$$S_8 + 8O_2 \xrightarrow{\Delta} 8SO_2$$

 $2MnO_4^- + 5SO_2 + 2H_2O \longrightarrow 5SO_4^{2-} + 4H^+ + 2Mn^{2+}$

$$2Fe^{3+} + SO_{2} + 2H_{2}O \longrightarrow 2Fe^{2+} + SO_{4}^{2-} + 4H^{+}$$

71.
$$Pb(NO_3)_2 \frac{\Delta}{673K} 2PbO + 4NO_2$$

(Brown colour)

$$2\mathrm{NO_2} \ \square \ \overset{\mathrm{On \ cooling}}{\mathrm{Heating}} \ \mathrm{N_2O_4}$$

(Colourless solid)

2NO +
$$\mathrm{N_2O_4}$$
 $_$ Δ 250 K \rightarrow 2 $\mathrm{N_2O_3}$

(Blue solid)

(Structure of N2O4)

$$\dot{\circ} = \dot{\circ} =$$

(Structure of N₂O₂)

72.
$$A = NH_4 NO_2$$
 $B = N_2$

 $C = NH_3$

 $D = HNO_3$

(i)
$$NH_4 NO_2 \rightarrow N_2 + 2H_2O$$

(ii)
$$N_2 + 3H_2 \rightarrow 2NH_3$$

(iii)
$$4\text{NH}_3 + 5\text{O}_2 \rightarrow 4\text{NO} + 6\text{H}_2\text{O}$$

 $4NO + O_2 \rightarrow 2NO_2$

 $3NO_2 + H_2O \rightarrow 2HNO_3 + NO$